

| | | |
|--|--|---|
| 15 | (?) <i>Magasella evansi</i> , Dall., <i>Terebratella evansi</i> , Dav., Proc. Zool. Soc., pl. xiv. fig. 7, 1852. See <i>Terebratella cruenta</i> . | New Zealand. |
| (?) | (?) <i>Magasella (Terebratula) rhombea</i> , Phil. Wieg. Archiv., 1845. Described, but not figured. Mr Dall tells me that he has seen the type at Berlin, and that it is identical with <i>Magasella evansi</i> (?). | Port Famine. |
| Genus <i>Laqueus</i> , Dall. | | |
| Loop long, three times attached to hinge-plate; again, by means of two horizontal processes, which branch off from the middle of principal stems to the mesial septum, and by slender processes from the reflected portion of loop to principal stems of loop. | | |
| 15 to 120 | <i>Laqueus californica</i> , Koch., nov. ed. Martini, vol. viii. p. 26, figs. 21-23. | Coasts of Sta Barbara, California; Port Etches, Prince William Sound to Catalina Island, Victoria. |
| 40 to 85 | <i>Laqueus picta</i> , Chemn., sp., Chemnitz, Conch. Tab., vol. xi. figs. 2011, 2012, 1785. Long considered a <i>Waldheimia</i> , but I am able to affirm that it belongs to the genus <i>Laqueus</i> . | Straits of Korea, 40 fathoms (Capt. St John); Japan, 50 to 85 fathoms. |
| 35 | <i>Laqueus rubella</i> , Sow., Thes. Conch., vol. i. p. 350, pl. lix. figs. 40-42, 1846 = <i>L. suffusa</i> , Dall, Am. Journ. Conch., vol. vi. p. 125, pl. vii. fig. 9, h, 1870. These are three good species. | Japan. |
| Genus <i>Megerlia</i> , King. | | |
| SECTION A.— <i>Megerlia</i> proper; loop three times attached, lateral loops closed, shell transverse, sculptured (Dall). | | |
| 10 to 292 | * <i>Megerlia truncata</i> , Lin., sp., <i>Syst. Nat.</i> , 1152, 1767 = <i>Anomia scobinata</i> , Gmelin = <i>T. decussata</i> and <i>T. irregularis</i> , De Blainville = <i>T. disculus</i> , Pallas = <i>Megerlia oblita</i> , Michelotti = <i>Morrisia gigantea</i> , Deshayes. | Mediterranean, Atlantic Shore; off Department of Finisterre (Collard des Chevres); off Groix, Morbihan; off Noir Mortiers, Vendée; and Cape Breton, France; off Guetaria, Spain (Hidaigo); Teneriffe 70 fathoms; off Island of Bourbon (?); Australia (?). <i>Fossil</i> : Upper Tertiaries; Italy; Nice; South of France; Austria. |
| 20 to 292 | <i>Megerlia truncata</i> , var. <i>monstruosa</i> , Scacchi, Osser. Zool., vol. ii. p. 1. This is considered by some malacologists to be a malformation of <i>M. truncata</i> . | Mediterranean. |
| 1850 | (?) * <i>Megerlia incerta</i> , Dav., Proc. Roy. Soc., vol. xxvii. p. 438, 1878. A rare deep-sea species; genus uncertain. | Between Sierra Leone and Fernando de Noronha. |