

Shore, tide mark.	<i>Kraussina davidsoni</i> , Vélain, Archiv. de Zool. experimentale, 1878. (The so-termed <i>K. picta</i> , alluded to by Mr Dall in his Catalogue of Recent Brachiopoda from the Island of St Paul, does not exist. The name is due to an erroneous reference in the Zool. Record.) <i>K. davidsoni</i> is so nearly related to <i>K. lamarckiana</i> , that it may, perhaps, be only a variety of that species.	Island of St Paul, Indian Ocean.
(?)	<i>Kraussina capensis</i> , Adams and Reeve, Voy. Samarang, Zool., p. 71, fig. 4.	Cape of Good Hope.
(?)	<i>Kraussina atkinsoni</i> , T. Wood's Census of Marine Shells of Tasmania, 1877. It is a small, smooth species.	Long Bay, Tasmania.
Genus <i>Bouchardia</i> , Dav.		
Apophysis, anchor-shaped; the septum being furnished with two short lamellæ.		
10 to 15	<i>Bouchardia rosea</i> , Mawe, sp., Introduction to Conchology, pl. xiv. fig. 4, 1823 = <i>T. tulipa</i> , Blainv. = <i>T. unguis</i> , Kuster.	Near Rio de Janeiro and Bahia.
Genus <i>Platydia</i> , Costa = <i>Morrisia</i> , Dav.		
Loop small, not reflected; attached to a small forked process in the centre of the valve.		
40 to 600	* <i>Platydia anomioides</i> , Scacchi, Moll. Sicil., vol. ii. p. 69, pl. xviii. fig. 9, 1844 = <i>Ter. appressa</i> , Forbes, 1844 = <i>semilunum</i> , Phil., according to Sequenza.	Mediterranean, 40 to 120 fms.; off Setubal, coast of Portugal, 600 fms. (Kent); off French Coast; Florida Reefs; off Marion and Prince Edward's Islands. <i>Fossil</i> : Upper Tertiaries; Sicily and South of Italy; Austria.
90 to 100	(?) <i>Platydia</i> (?) <i>davidsoni</i> , E. Desl., Ann. and Mag. of Nat. Hist., 2d ser., vol. x. fig. 20, 1855; and P. Fischer, Journ. de Conch., 3d ser., vol. xii. p. 160, 1872. Dr Gwyn Jeffreys considers <i>P. davidsoni</i> to be the same as <i>Megerlia monstruosa</i> (?), but this opinion will require confirmation, as both Mr E. Deslongchamps and P. Fischer are of a contrary opinion.	Mediterranean, off Corsica; Coral Fisheries, near Tunis; off Cape Breton, France (Fischer). <i>Fossil</i> : Upper Tertiaries, Sicily.
59 to 95	(?) <i>Platydia</i> (?) <i>lunifera</i> , Phil., Moll. Sicil., vol. i. p. 97, pl. vi. fig. 16, 1836. This species and the genus to which it belongs are still uncertain.	Mediterranean.
Genus <i>Argiope</i> , E. Deslongchamps.		
Loop four-lobed, adhering to three or five sub-marginal septa.		
18 to 364	<i>Argiope decollata</i> , Chemnitz, Conch. Cab., vol. viii. p. 96, pl. lxxviii., fig. 705 = <i>A. detruncata</i> , Gmel. = <i>T. ungula</i> , Retz = <i>T. aperta</i> , Blainv. = <i>T. pectiniformis</i> , Costa =	Two miles east of Guernsey; Atlantic Coast of France; off Cape Breton; Mediter-