

690	<i>Rhynchonella sicula</i> , Sequenza. See Jeffreys, Proc. Zool., 1878, p. 413, pl. xxiii. figs. 5, 6. It seems to be a good and well characterised species, more specimens required.	Chops of English Channel, "Porcupine" Expedition (Jeffreys). <i>Fossil</i> : South of Italy; Sicily.
15 to 150	<i>Rhynchonella nigricans</i> , Sow., Thes. Conch. vol. i. p. 342, pl. lxxi. fig. 81 82, 1846. A very good and remarkable species.	Fouveaux Strait, New Zealand; Chatham Islands.
150	* <i>Rhynchonella nigricans</i> , var. <i>picylata</i> , R. B. Watson, MS., Dav., Proc. Roy. Soc., vol. xxvii. p. 431, 1878.	South of Kerguelen Islands. <i>Fossil</i> (?): Tasmania.
(?)	<i>Rhynchonella grayi</i> , Woodward, Ann. and Mag. of Nat. Hist., p. 444, pl. x. fig. 16, 1855. Only one example known. British Museum. More specimens required.	Off Fiji Islands.
48 to 100	<i>Rhynchonella lucida</i> , Gould, Proc. Boston Soc. Nat. Hist., p. 323, 1860. Seems to be a very good and well-characterised species.	Japan.
TRETENTERATA.		
Genus <i>Crania</i> , Retzius.		
3 to 808	<i>Crania anomala</i> , Müller, Zool. Dan., p. 237, 1776 = <i>Patella distorta</i> , Montagu. <i>Patella kermes</i> , Humphreys = <i>Anomia turbinata</i> , Dillwyn = <i>Criopus orcadensis</i> , Leich = <i>Orbicula norvegica</i> , Lam. <i>D. ostreoides</i> , Turton = <i>C. personata</i> , Def. = <i>C. rostrata</i> , Thorpe. <i>C. fimbriata</i> , Poli = <i>C. craniolaris</i> (pars), Gmel. = <i>C. ringens</i> , Høninghaus.	North European Seas; Loch Fyne; North of Scotland; Tunisian Coast, Mediterranean, from 14 to 120 fathoms; off Province of Asturias, Spain; Vigo (M'Andrew); off Biarritz (Fischer). <i>Fossil</i> : Sicily (?).
(?)	<i>Crania anomala</i> , var. <i>alba</i> , Jeffreys, Brit. Conch., vol. v. p. 165, 1869. I am not acquainted with this or the following variety.	Off Shetland Islands; Hebrides.
(?)	<i>Crania anomala</i> , var. <i>pourtalesi</i> , Dall, Bull. Mus. Comp. Zool., vol. i. p. 35, pl. i. fig. 7, 1871.	Florida Keys.
71	<i>Crania japonica</i> , A. Adams, Ann. and Mag. Nat. Hist., 3d ser., vol. xi. p. 100, 1863.	Japan.
Low water.	<i>Crania suessi</i> , Reeve, Conch. con. Mon. <i>Crania</i> , pl. i. fig. 2, 1862. Both these last seem to be good species.	Sydney, Australia.
Genus <i>Discina</i> , Lamarck.		
Mr Dall proposes to divide the genus <i>Discina</i> as follows:—		
A. <i>Discina</i> , Lamarck, with anteriorly directed tubular foramen, under the central septum or prominence (Dall).		
6 to (?)	<i>Discina striata</i> (<i>Crania</i>), Schumacher, Essai, p. 102, pl. xx. fig. 1, a-f, 1817 (not of DeFrance) = <i>C. radiosa</i> , Gould = <i>D. evansi</i> , Dav., <i>Orbicula norvegica</i> , Sow. <i>O. ostreoides</i> , Rang. <i>Patella anomala</i> , Sow. (not Müller).	North-West coast of Africa.