

	Range in Depth. Fathoms.	Principal Localities.
<i>Echinothrix</i> , Pet.		
× <i>E. calamaris</i> , A. Ag.,	—	{ Society Islands; East India Islands; Philippine Islands; Kandavu Reef. Red Sea; Fiji Islands; Sandwich Islands. Sandwich Islands; Fiji Islands; Japan; East India Islands; Red Sea; Zanzibar; Kandavu Reef.
<i>E. desorii</i> , Pet.,	—	
× <i>E. turcarum</i> , Pet.,	—	
<i>Astropyga</i> , Gray.		
<i>A. elastica</i> , Stud.,	[20	New Britain. Panama; Gulf of California; Torres Straits; Honolulu. Zanzibar; East India Islands; Philippine Islands.
× <i>A. pulvinata</i> , Agass.,	-49 C.	
<i>A. radiata</i> , Gray,	—	
ECHINOTHURIDÆ, Wy. Thom.		
<i>Asthenosoma</i> , Grube.		
* <i>A. grubii</i> , A. Ag.,	[10 C.	Philippines. China Seas.
<i>A. varium</i> , Grube,	—	
ECHINOMETRADÆ, Gray.		
<i>Colobocentrotus</i> , Br.		
<i>C. atratus</i> , Br.,	—	Zanzibar; Java; Sandwich Islands. Bonin Islands; Australia.
<i>C. mertensii</i> , Br.,	—	
<i>Heterocentrotus</i> , Br.		
× <i>H. mammillatus</i> , Br.,	—	{ Sandwich Islands; East India Islands; Red Sea; Fiji Islands; Kandavu Reef. Maritius; Java; Sandwich Islands; Fiji Islands; New Caledonia.
× <i>H. trigonarius</i> , Br.,	—	
<i>Echinometra</i> , Rondel. (Breyn.)		
× <i>E. lucunter</i> , Bl.,	-18 C.	{ Zanzibar; Red Sea; East India Islands; Japan; Sandwich Islands; Fiji Islands; Philippines; Kandavu Reef. West Coast, Africa. Sandwich Islands; Philippine Islands; Seychelle Islands. Senegal; Cape Verde; Brazil; West India Islands; Ber- mudas; Ascension. Peru; Panama; Gulf of California. Florida; Hayti.
<i>E. macrostoma</i> , A. Ag.,	—	
<i>E. oblonga</i> , Bl.,	—	
× <i>E. subangularis</i> , Desml.,	-250 B.	
<i>E. van brunti</i> , A. Ag.,	—	
<i>E. viridis</i> , A. Ag.,	-7 B.	
<i>Parasalenia</i> , A. Ag.		
<i>P. gratiosa</i> , A. Ag.,	—	{ Kingsmills Islands; Bonin Islands; Zanzibar; Fiji Islands; New Caledonia.
<i>Stomopneustes</i> , Agass.		
<i>S. variolaris</i> , Agass.,	—	Mauritius; Java; Samoa.
<i>Strongylocentrotus</i> , Br.		
<i>S. albus</i> , A. Ag.,	—	Patagonia; Chili; Peru; Philippines. Australia.
<i>S. armiger</i> , A. Ag.,	—	
<i>S. depressus</i> , A. Ag.,	—	Japan.
× <i>S. dröbachiensis</i> , A. Ag.,	-78	{ North European; North Pacific; North-East Coast of America. Australia; Tasmania; Samoa; New Caledonia; New South Wales.
× <i>S. eurythrogrammus</i> , A. Ag.,	-35 C.	
<i>S. franciscanus</i> , A. Ag.,	—	Formosa; Puget Sound; San Diego; Japan.
<i>S. gaimardi</i> , A. Ag.,	—	Brazil.
× <i>S. gibbosus</i> , A. Ag.,	-45 C.	Chili; Galapagos; Peru; Fiji Islands; West Patagonia. Japan; Australia.
<i>S. intermedius</i> , A. Ag.,	—	
<i>S. lividus</i> , Br.,	—	European Atlantic; Mediterranean; Azores.
<i>S. mexicanus</i> , A. Ag.,	—	Gulf of California.
<i>S. nudus</i> , A. Ag.,	—	Sandwich Islands; Japan.
<i>S. purpuratus</i> , A. Ag.,	—	San Francisco; Puget Sound.
<i>S. tuberculatus</i> , Br.,	—	Japan; China; Australia; New Zealand; Galapagos.