

	Range in Depth and Principal Localities.
<i>Echinus</i> , Rond. (Linn.)	
⊕ <i>E. acutus</i> , Lamk.,	× Station 343, 425 fathoms (Atlantic; Pacific).
⊕ <i>E. elegans</i> , Düb. o. Kor.,	× Tristan da Cunha, 1100 fathoms (Atlantic; Pacific).
<i>E. gracilis</i> , A. Ag.,	{ Florida, Gulf Stream, 100-200 fathoms; St Thomas; Caribbean, 175 fathoms B.
<i>E. norvegicus</i> , Düb. o. Kor.,	{ Florida, Gulf Stream, 195 fathoms; Atlantic Coast of U. S., 1242 fathoms B.; (Atlantic; Japan).
<i>E. wallisii</i> , A. Ag.,	Atlantic Coast of U. S. (South and Middle States), 267-1047 fathoms B.
<i>Toxopneustes</i> , Agass.	
× <i>T. variegatus</i> , A. Ag.,	{ North Carolina, 7 fathoms; Bermudas; Florida, Gulf Stream, 5-34 fathoms; Caribbean Islands, 14-242 fathoms B.
<i>Hipponoë</i> , Gray.	
<i>H. esculenta</i> , A. Ag.,	{ South Florida; Bermudas; Surinam; Caribbean Islands, 14-451 fathoms B.
CLYPEASTRIDÆ, Agass.	
ECHINONIDÆ, D'Orb.	
<i>Pygaster</i> , Agass.	
<i>P. relictus</i> , Lov.,	Virgin Islands, 180 fathoms J.
EUCLYPEASTRIDÆ, Hæckel.	
FIBULARINA, Gray.	
<i>Echinocyamus</i> , Van Phelps.	
× <i>E. pusillus</i> , Gray,	{ Florida, Gulf Stream, 5-325 fathoms; Caribbean Islands, 98-805 B.; × Station 122. Off Barra Granda (no depth on label), (North Atlantic).
ECHINANTHIDÆ, A. Ag.	
<i>Clypeaster</i> , Lamk.	
× <i>C. subdepressus</i> , Agass.,	{ South Carolina; Brazil; Florida, Gulf Stream, 11-40 fathoms; Florida; Caribbean Islands, 36-248 fathoms B. (North Atlantic).
<i>Echinanthus</i> , Breyn.	
<i>E. rosaceus</i> , Gray,	South Carolina; West Indies; Caribbean Islands, 14-118 fathoms B.
SCUTELLIDÆ, Agass.	
<i>Mellita</i> , Kl.	
× <i>M. sexforis</i> , A. Ag.,	{ South Carolina; Bermudas; West Indies; Florida, Gulf Stream, 4-270 fathoms; × Bahia.
<i>M. testudinata</i> , Kl.,	East Virginia; West Indies; Rio Janeiro.
<i>Encope</i> , Agass.	
× <i>E. emarginata</i> , Agass.,	{ South Carolina; West Indies; Florida, Gulf Stream, 7 fathoms; Rio Janeiro; × Bahia, 20-70 fathoms.
<i>E. michelini</i> , Agass.,	{ Alabama; West Florida; Honduras; Yucatan; Florida, Gulf Stream, 5-6 fathoms; Caribbean Islands, 14 fathoms B.
PETALOSTICHA, Hæckel.	
CASSIDULIDÆ, Agass.	
ECHINONIDÆ, Agass.	
<i>Echinonëus</i> , Van Phelps.	
<i>E. semilunaris</i> , Lamk.,	Bermudas; Florida Keys; Caribbean Islands, 80-805 fathoms B.
NUCLEOLIDÆ, Agass.	
<i>Neolampas</i> , A. Ag.	
<i>N. rostellata</i> , A. Ag.,	{ Florida, Gulf Stream, 112-125 fathoms; Caribbean Islands, 100-229 fathoms B. (North Atlantic).