

	Range in Depth and Principal Localities.
ECHINANTHIDÆ, A. Ag.	
<i>Clypeaster</i> , Lamk. <i>C. subdepressus</i> , A. Ag., . . .	West Coast of Tropical Africa (North Atlantic).
SCUTELLIDÆ, Agass.	
<i>Rotula</i> , Kl. <i>R. augusti</i> , Kl., : : : : <i>R. rumpfii</i> , Kl., : : : :	West Africa; Liberia. Cape Verde Islands; Porto Praya, 20 fathoms; Senegal.
PETALOSTICHA , Hæckel.	
CASSIDULIDÆ, Agass.	
NUCLEOLIDÆ, Agass.	
<i>Echinolampas</i> , Gray. <i>E. hellei</i> , Val.,	Liberia; Senegal.
SPATANGIDÆ, Agass.	
POURTALESIDÆ, A. Ag.	
<i>Calymene</i> , Wy. Thom. * <i>C. relicta</i> , Wy. Thom.,	*Fayal, 2650 fathoms.
ANANCHYTIDÆ, Alb. Gras.	
<i>Palaeotropus</i> , Lov. <i>P. josephinae</i> , Lov.,	Azores, 250 fathoms, J. (North Atlantic).
Homolampas, A. Ag.	
<i>H. fragilis</i> , A. Ag.,	Josephine Bank, 5–600 fathoms, J. (North Atlantic).
SPATANGINA, Gray.	
<i>Spatangus</i> , Kl. × <i>S. purpureus</i> , Leske, <i>S. raschi</i> , Lov.,	{ Mediterranean; Adriatic; × Azores, × Station 75, 50–90 fathoms (North Atlantic). Azores; Faroe Islands—Gibraltar, 100–300 fathoms, P. (Atlantic).
Echinocardium, Gray.	
<i>E. cordatum</i> , Gray, <i>E. mediterraneum</i> , Gray,	Mediterranean (North Atlantic). Mediterranean; Sicily; Messina, 2–20 fathoms.
BRISSINA, Gray.	
Hemaster , Des. <i>H. expurgatus</i> , Lov., * <i>H. zonatus</i> , A. Ag.,	West Coast of Spain, 485 fathoms, J. (North Atlantic). × Station VIII., 620 fathoms (North Atlantic).
Brissopsis , Agass. <i>B. lyra</i> , Agass.,	Mediterranean; Sicily (Atlantic).
Aerope , Wy. Thom. * <i>A. rostrata</i> , Wy. Thom.,	× Bay of Biscay and Coast of Portugal (Atlantic; Pacific).
Acoste , Wy. Thom. * <i>A. bellidifera</i> , Wy. Thom.,	× Station VIII., 620 fathoms (Atlantic; Pacific).