

	Range in Depth and Principal Localities.
<i>Brissus</i> , Kl.	
* <i>B. dumesi</i> , A. Ag., . . . .	×Station 75, 450 fathoms (North Atlantic).
<i>B. unicolor</i> , Kl., . . . .	Cape Verde Islands; Sicily (North Atlantic).
<i>Metalia</i> , Gray.	
<i>M. africana</i> , A. Ag., . . . .	West Coast Tropical Africa.
<i>M. costæ</i> , Ludl., . . . .	Capri, 25 fathoms.
<i>Schizaster</i> , Agass.	
<i>S. canaliferus</i> , Agass., . . . .	Mediterranean, 37 fathoms; Sicily.

## CAPE OF GOOD HOPE—NATAL.

	Range in Depth and Principal Localities.
DESMOSTICHA, Hæckel.	
CIDARIDÆ, Müll.	
GONIOCIDARIDÆ, Hæckel.	
<i>Cidaris</i> , Kl.	
<i>C. metularia</i> , Bl., . . . .	(African-Indian-Pacific).
<i>Phyllacanthus</i> , Br.	
<i>P. baculosa</i> , A. Ag., . . . .	(Indo-African).
<i>P. dubia</i> , Br., . . . .	(Indo-African).
<i>Goniocidaris</i> , Des.	
<i>G. canaliculata</i> , A. Ag., . . . .	(Southern Ocean).
ARBACIADÆ, Gray.	
<i>Arbacia</i> , Gray.	
× <i>A. dufresnii</i> , Gray, . . . .	×Nightingale Island (Southern Ocean).
DIADEMATIDÆ, Peters.	
<i>Diadema</i> , Schynv.	
<i>D. setosum</i> , Gray, . . . .	(Atlantic; Pacific).
<i>Echinothrix</i> , Pet.	
<i>E. desorii</i> , Pet., . . . .	(African-Indian-Pacific).
ECHINOMETRADÆ, Gray.	
<i>Echinometra</i> , Rond. (Breyn.)	
<i>E. lucunter</i> , Bl., . . . .	Natal (African-Indian-Pacific).
<i>Stomopneustes</i> , Agass.	
<i>S. variolaris</i> , A. Ag., . . . .	Natal (Indo-African).
<i>Echinostrephus</i> , A. Ag.	
<i>E. molare</i> , A. Ag., . . . .	Natal (African-Indian-Pacific).
ECHINIDÆ, Agass.	
TRIPLECHINIDÆ, A. Ag.	
<i>Echinus</i> , Rond. (Linn.)	
× <i>E. angulosus</i> , A. Ag., . . . .	×Simon's Bay, 10-20 fathoms (African-Indian-Southern Ocean).