

|   | Range in Depth and Principal Localities.  |
|---|---|
| SCUTELLIDÆ, Agass.                      |   |
| <i>Echinarachnius</i> , Leske.          |   |
| <i>E. mirabilis</i> , A. Ag., . . . .   | Hakodadi.   |
| <i>Echinodiscus</i> , Breyn.            |   |
| <i>E. lœvis</i> , A. Ag., . . . .       | China Sea, 20 fathoms; Ousima (Indo-African).   |
| <i>Astrichlypeus</i> , Verrill.         |   |
| × <i>A. manni</i> , Verrill, . . . .    | China; Yeddo; × Inland Sea, Japan.  |
| PETALOSTICHA, Hæckel.                   |   |
| SPATANGIDÆ, Agass.                      |   |
| POURTALESIÆ, A. Ag.                     |   |
| <i>Pourtalesia</i> , A. Ag.             |   |
| * <i>P. laguncula</i> , A. Ag., . . . . | × Station 232, 345 fathoms (Pacific).   |
| <i>Cystechinus</i> , A. Ag.             |   |
| * <i>C. clypeatus</i> , A. Ag., . . . . | × Station 205, 1050 fathoms (South Atlantic).   |
| ANANCHYTIDÆ, Alb. Gras.                 |   |
| <i>Linopneustes</i> , A. Ag.            |   |
| * <i>L. murrayi</i> , A. Ag., . . . .   | × Station 232, 345 fathoms (Pacific).   |
| SPATANGINA, Gray.                       |   |
| <i>Spatangus</i> , Kl.                  |   |
| <i>S. lilkeni</i> , A. Ag., . . . .     | Formosa; Hakodadi.  |
| <i>Maretia</i> , Gray.                  |   |
| <i>M. alta</i> , A. Ag., . . . .        | Kagosima (Indian).  |
| <i>M. planulata</i> , Gray, . . . .     | China (African-Indian-Pacific).   |
| <i>Lovenia</i> , Des.                   |   |
| × <i>L. subcarinata</i> , Gray, . . . . | China; Hakodadi; × Hong Kong, 10 fathoms (Indian-Pacific).  |
| <i>Echinocardium</i> , Gray.            |   |
| × <i>E. australe</i> , Gray, . . . .    | { China; Japan; × Kobi, 8-50 fathoms; × Station 234, 2675 fathoms (Indian-Southern Ocean).                                |
| LFSKIADÆ, Gray.                         |   |
| <i>Palæostoma</i> , Lovén.              |   |
| <i>P. mirabilis</i> , Lov., . . . .     | Hong Kong (Indian).   |
| BRISSINA, Gray.                         |   |
| <i>Hemiaster</i> , Des.                 |   |
| * <i>H. gibbosus</i> , A. Ag., . . . .  | × Station 232, 345 fathoms (Pacific).   |
| <i>Rhinobriassus</i> , A. Ag.           |   |
| <i>R. pyramidalis</i> , A. Ag., . . . . | China.  |
| <i>Brissopsis</i> , Agass.              |   |
| × <i>B. luzonica</i> , A. Ag., . . . .  | { Formosa; × Station 203, 12-20 fathoms; × Station 232, 345 fathoms (Indo-Pacific).                                       |
| <i>Faorina</i> , Gray.                  |   |
| <i>F. chinensis</i> , Gray, . . . .     | China; Hong Kong; Shanghai (Pacific?).  |
| <i>Schizaster</i> , Agass.              |   |
| * <i>S. japonicus</i> , A. Ag., . . . . | { × Yokohama, 8-14 fathoms; × Kobi, Japan, 8-50 fathoms; × Hong Kong, 10 fathoms; × Station 233 B., 15 fathoms (Pacific). |
| × <i>S. ventricosus</i> , Gray, . . . . | Hong Kong; × Station 232, 345 fathoms (Indo-Pacific).   |