

Of the Japanese and Chinese species *Schizaster japonicus* is the only new littoral species collected by the Challenger. The habitat of *Pleurechinus* is determined for the first time, and *Echinus norvegicus* and *Dorocidaris papillata* are recorded for the first time as coming from the Pacific. As far as the other deep-water species are concerned there are of the continental Indo-African *Cælopleurus maillardii*, *Micropyga tuberculatum*, while *Podocidaris*, *Phormosoma*, *Asthenosoma*, and *Pourtalesia* are Pacific abyssal, with the exception of *Asthenosoma varium*, and the Japanese representative continental *Linopneustes murrayi* and *Goniocidaris florigera*.

PACIFIC OCEAN.

	Range in Depth and Principal Localities.
DESMOSTICHA , Hæckel. CIDARIDÆ , Müll. GONIOLIDARIDÆ , Hæckel.	
<i>Cidaris</i> , Kl <i>C. metularia</i> , Bl,	Solomon and Sandwich Islands (African-Indian-Pacific).
<i>Dorocidaris</i> , A. Ag. * <i>D. bracteata</i> , A. Ag., ⊕ <i>D. papillata</i> , A. Ag.,	× Amboyna, 15–100 fathoms. × Station 210, 375 fathoms (Atlantic).
<i>Phyllacanthus</i> , Br. × <i>P. annulifera</i> , A. Ag., × <i>P. baculosa</i> , A. Ag., <i>P. dubia</i> , Br., × <i>P. gigantea</i> , A. Ag., <i>P. imperialis</i> , Br., × <i>P. verticillata</i> , A. Ag.,	{ South Sea; × Cape York; × Station 186, 8 fathoms; × Station 188, 28 fathoms (Indian). Samboangan, 10 fathoms (Indo-African). Bonin Islands; New Caledonia (Indo-African). Sandwich Islands (African-Indian-Pacific). Tonga (African-Indian-Pacific). { Navigator Islands; Society Islands; Sandwich Islands; × Station 186, 8 fathoms (African-Indian-Pacific).
<i>Porocidaris</i> , Des. * <i>P. elegans</i> , A. Ag.,	× Station 214, 500 fathoms.
<i>Goniocidaris</i> , Des. <i>G. canaliculata</i> , A. Ag., * <i>G. florigera</i> , A. Ag.,	Caroline and Sandwich Islands (Southern Ocean). × Station 192, 129 fathoms.
SALENIDÆ , Agass.	
<i>Salenia</i> , Gray. * <i>S. hastigera</i> , A. Ag.,	{ × Station 170, 630 fathoms; × Station 171, 600 fathoms; × Station 195, 1425 fathoms; × Off Cebu, 100 fathoms (Atlantic).
ARBACIADÆ , Gray.	
<i>Podocidaris</i> , A. Ag. * <i>P. prionigera</i> , A. Ag.,	× Station 218, 1070 fathoms.
<i>Cælopleurus</i> , Agass. ⊕ <i>C. maillardii</i> , A. Ag.,	× Station 192, 129 fathoms; × Amboyna, 100 fathoms (Indo-African).