

Ophiacantha aspera, Lym., Bull. Mus. Comp. Zool., vol. v., part 9, p. 228, pl. i. figs. 10-12, 1878.

Off Havana; 175 fathoms.

Ophiacantha indica, Ljn., Oph. Viv., Öf. Kong. Akad., p. 326, 1866.

Between Batavia and Singapore.

Ophiacantha dallasii, Duncan, Linn. Soc. Journ., vol. xiv. p. 471, pl. xi. figs. 25-27, 1879.

38° 19' N. 129° 7' E.; 50 fathoms.

Ophiacantha stellata, Lym., Ill. Cat. Mus. Comp. Zool., No. viii., part 2, p. 11, pl. ii. figs. 16-18, 1875.

Barbadoes; 100 fathoms.

Ophiacantha scutata, Lym., Bull. Mus. Comp. Zool., vol. v., part 9, p. 229, pl. i. figs. 1-3, 1878.

West Indies; 287 to 292 fathoms.

Ophiacantha pentacrinus, Ltk., Addit. ad Hist., part 3, pp. 46 and 99, 1869; Lym., Bull. Mus. Comp. Zool., vol. v., part 9, p. 230.

Ophiacantha meridionalis, Lym., Bull. Mus. Comp. Zool., vol. i., part 10, p. 324, 1869.

West Indies; 74 to 539 fathoms.

Ophiacantha abyssicola, G. O. Sars, Nye Echin. Vid. Selsk. Forh., p. 8, 1871.

Lofoten Island; 20 to 300 fathoms.

Ophiacantha marsupialis, Lym., Ill. Cat. Mus. Comp. Zool., No. viii., part 2, p. 13, pl. i. figs. 9, 10, 1875.

Juan Fernandez; 240 fathoms.

Ophiolebes.

Ophiolebes, Lym., Bull. Mus. Comp. Zool., vol. v., part 7, 1878.

Disk and arms stout, and covered by a thick skin, which bears grains or stumps, and hides more or less the underlying plates. Arm spines short, blunt, thorny, covered by thick skin, and arranged on the sides of the side arm plates, which project but slightly. Long, stout mouth papillæ, and teeth; no tooth papillæ; peristomial plates large and wide, making a circle by their connecting ends. Arm plates rather small, but normal, nearly as in *Ophiacantha*. Two genital openings in each interbrachial space.