

Table of Species of *Ophiothrix*—continued.

Arms five to six times diameter of disk. Radial shields essentially naked. A slight re-entering curve in outer side of under arm plate.	Disk beset with minute trifid stumps, among which a few slender spines. Colour in alcohol, blue, or pale pink, with a light line along upper arm,	} <i>Ophiothrix angulata</i> .
	Disk beset with slender spines. Colour in alcohol, blue or green, with fine white cross-lines on arms,	} <i>Ophiothrix ørstedii</i> .
	Disk beset with thorny stumps and slender spines, or with either alone. Arm spines longer than in <i>Ophiothrix angulata</i> . Colour in alcohol, pale blue,	} <i>Ophiothrix spiculata</i> .
	A few slender spines on disk. Several thin lines alternately blue and white, running along upper arm,	} <i>Ophiothrix trilineata</i> .
	Similar to <i>Ophiothrix spiculata</i> , but under arm plates proportionately longer; and the lowest arm spine keeps the form of a double hook till quite near the base of the arm,	} <i>Ophiothrix koreana</i> .
	Disk beset with coarse spines of several sizes. Arm spines stouter than in <i>Ophiothrix spiculata</i> , and under arm plates more angular,	} <i>Ophiothrix dumosa</i> .
	Large (disk 14 mm.). Arm spines rounded, stout, and tapering. Disk closely set with similar smaller spines,	} <i>Ophiothrix magnifica</i> .
Some of the arm spines clubbed or widened at the end. Radial shields naked.	Disk with a few peg-like stumps. Upper arm plates hexagonal. Arm spines very stout. Colour in alcohol reddish, with black line along upper arm,	} <i>Ophiothrix lineata</i> .
	Radial shields small and set, like disk, with small thorny stumps. Under arm plates with a re-entering curve on outer side. Arm spines thick and not strongly thorned,	} <i>Ophiothrix triglochis</i> .
	Disk as in <i>Ophiothrix triglochis</i> , but arm spines much flatter and more toothed; and under arm plates are curved outwardly,	} <i>Ophiothrix caespitosa</i> .
	Radial shields naked. Upper disk set with minute, smooth, cylinders. Seven nearly cylindrical, tapering arm spines, only a little thorny at the tip. Under arm plates covered without,	} <i>Ophiothrix rudis</i> .
	Radial shields and lower interbrachial spaces nearly naked; rest of disk densely set with stumps bearing a crown of thorns. Outer side of under arm plates curved. Seven short, flat, strongly toothed arm spines,	} <i>Ophiothrix berberis</i> .
	Radial shields naked. Upper arm spines club-ended. Arm to disk 7:1. Short spines on disk, which is spotted with black, and arms banded with same,	} <i>Ophiothrix spongicola</i> .
	Radial shields naked. A few spines on disk. Arm spines with shaft smooth and a flat thorny end. Lower arm plates as broad as long and curved without,	} <i>Ophiothrix melanosticta</i> .
	Disk as in preceding. Arm spines with smooth shaft and rough end; the second with a broadened flat, toothed end. Under arm plates squarish with an outer re-entering curve,	} <i>Ophiothrix striolata</i> .
	Disk naked above, spinous below. Upper arm spines with smooth cylindric shaft and thorny clubbed end. Lowest one a trifle hooked. Arm to disk 4½:1,	} <i>Ophiothrix martensi</i> .
	Disk as in <i>Ophiothrix martensi</i> . Outer side of under arm plate re-entering curved. Most arm spines with brush like thorny ends. Arms to disk 5:1,	} <i>Ophiothrix plana</i> .