

PLATE XXX.

- Fig. 1. *Astroschema horridum*, Lym., below, with one arm complete, $\frac{4}{1}$.
 „ 2. „ „ above, $\frac{4}{1}$.
 „ 3. „ „ arm joint near base of arm, $\frac{4}{1}$.
 „ 4. „ „ arm joint some distance along the arm, $\frac{8}{1}$.
 „ 5. „ *brachiatum*, Lym., below, $\frac{6}{1}$.
 „ 6. „ „ above with one arm complete, $\frac{6}{1}$.
 „ 7. „ „ arm joint near base of arm in profile, $\frac{6}{1}$.
 „ 8. „ „ arm joint some distance along the arm in profile, $\frac{6}{1}$.

PLATE XXXI.

(The alcoholic specimens were allowed partially to dry, and the outlines are, therefore, too sharp.)

- Fig. 1. *Ophiocreas carnosus*, Lym., below, $\frac{2}{1}$.
 „ 2. „ „ above, $\frac{2}{1}$. (The disk should be more wrinkled and fleshy.)
 „ 3. „ „ arm joint near base of arm, $\frac{3}{2}$.
 „ 4. „ „ arm joint near tip of arm, $\frac{2}{1}$.
 „ 5. „ *adipus*, Lym., below, $\frac{4}{1}$.
 „ 6. „ „ above, $\frac{4}{1}$.
 „ 7. „ „ arm joint near base of arm, $\frac{6}{1}$.
 „ 8. „ „ arm joint near tip of arm, $\frac{4}{1}$.

PLATE XXXII.

(The alcoholic specimens were allowed partially to dry, and the outlines are, therefore, too sharp.)

- Fig. 1. *Ophiocreas abyssicola*, Lym., above, $\frac{4}{1}$. (The radial shields are too strongly marked, and they do not go to the centre.)
 „ 2. „ „ below, $\frac{4}{1}$. (The mouth angle is incorrect; there are no mouth papillæ, only irregular grains.)
 „ 3. „ „ arm joint near base of arm, $\frac{5}{1}$.
 „ 4. „ „ arm joint near tip of arm, $\frac{5}{1}$.
 „ 5. „ *caudatus*, Lym., below, $\frac{2}{1}$.
 „ 6. „ „ above, $\frac{2}{1}$.
 „ 7. „ „ arm joint near base of arm, $\frac{2}{1}$.
 „ 8. „ „ arm joint near tip of arm, $\frac{5}{1}$.