## PLATE XII.

- Figs. 1-7. Culeolus recumbens, Herdman. Figs. 8-12. Culeolus perlucidus, Herdman.
- Fig. 1. A section through the test of Culeolus recumbens, showing foreign particles imbedded in the outer part (left side of figure); magnified (Hartn., obj. 4).
- Fig. 2. A transverse section through the peduncle of the same species, showing the cavities (ped.c.), and the imbedded foreign particles; magnified (Hartn., obj. 4).
- Fig. 3. Part of the branchial sac, from the inside, showing a fold (br. f.); magnified (Hartn., obj. 4).
- Fig. 4. Part of the endostyle, showing the clear edges, with spicules, and the central opaque area; magnified (Hartn., obj. 4).
- Fig. 5. Columnar cells from the brown band of the endostyle; much magnified (Hartn., obj. 7).
- Fig. 6. The terminations of the genital ducts; magnified (Hartn., obj. 4).
- Fig. 7. One of the genital glands, enlarged, showing the ducts.
- Fig. 8. A section through the test of *Culeolus perlucidus*, showing the papillæ on the outer surface; magnified (Hartn., obj. 4).
- Fig. 9. Part of the endostyle, showing the clear edge (w.e.), and the central more opaque area (c.b.a.); magnified (Hartn., obj. 4).
- Fig. 10. Part of the circlet of tentacles, showing three orders (tn., tn'. and tn."); magnified (Hartn., obj. 4).
- Fig. 11. Part of a section through one of the genital glands, showing the spermatic vesicles externally, and the ova internally; magnified (Hartn., obj. 4).
- Fig. 12. Part of the wall of the stomach, with the lining epithelium scraped off, showing the branched hepatic (?) tubules, with enlarged ends; much magnified (Hartn., obj. 7).