

23	<i>L. variata</i> , Brady. Often unsymmetrical or depressed; with irregular, depressed striæ,	Pl. LXI. fig. 1.
24	Lagena sulcata , W. & J., sp. Surface ornamented with stout raised longitudinal costæ,	Pl. LVII. figs. 23, 33, &c.
25	<i>interrupta</i> , Will. Costæ of unequal length,	— figs. 25, 27.
26	<i>acuticosta</i> , Reuss, with few, thin, much elevated costæ; sometimes irregularly placed,	— figs. 31, 32.
27	<i>diversicostata</i> , Reuss. With few costæ, divided into two sets by a deep depression; the two central costæ continuous across the base of the test,	G.
28	<i>williamsoni</i> , Alcock, sp. Pyriform, costate, with hexagonal ornament round the base of the neck,	H.
29	<i>L. gracilis</i> , Will. Elongate, fusiform, costate or striate; pointed at the inferior end; sometimes distomous,	Pl. LVIII. figs. 2, 3, 7, 8.
30	<i>L. alifera</i> , Reuss. Costæ very thin and high, taking the form of wings,	I.
31	<i>L. quadralata</i> , Brady. Flask-shaped, finely striate; with four equidistant wings,	Pl. LXI. fig. 3.
32	<i>L. plumigera</i> , Brady. Flask-shaped; with costæ developed at their lower ends into broad tubulated wings,	Pl. LVIII. figs. 25-27.
33	Lagena semistriata , Williamson. Inferior portion of the test costate, superior portion smooth,	Pl. LVII. figs. 14, 16, &c.
34	<i>tricostulata</i> , Marsson. Globular; with three parallel costæ reaching to the middle of the test,	J.
35	<i>multicosta</i> , Karrer, sp. Subglobular; aperture fissurine; with simple or bifurcating costæ reaching to the middle of the test,	Pl. LXI. fig. 4.
36	<i>L. stelligera</i> , Brady. Pyriform; with short costæ proceeding from a raised rim or collar at the base,	Pl. LVII. figs. 35, 36.
37	<i>L. lebouriana</i> , Brady. Inferior end costate; the ribs connected by a transverse raised band encircling the test near the middle,	K.
38	<i>L. gothica</i> , Ry. Jones. Lower half costate; the superior ends of the adjacent costæ curved over so as to form gothic arches,	L.
39	<i>L. crenata</i> , P. & J. Decanter-shaped; the broad base ornamented with radiate crenulations proceeding from a small convex central disk,	— fig. 21.
40	<i>L. exsculpta</i> , Brady. Ornament consisting of an excavated star, extending to near the middle of the test; radii fluted, bordered, round at the top,	Pl. LVIII. fig. 1, &c.
41	Lagena striatopunctata , Parker & Jones. Multicostate; with conspicuous perforations at regular intervals on the costæ,	— figs. 37, 40.
42	<i>howchiniana</i> , Brady. Test crenulated or sub-costate; ridges perforated; wall of even thickness,	M.
43	<i>tetragona</i> , P. & J. Quadricostate, striato-punctate,	N.
44	<i>spiralis</i> , Brady. Striato-punctate, costæ spiral,	Pl. CXIV. fig. 9.