

PLATE XLI.

TROCHAMMINA.—CARTERINA.—WEBBINA.

		Diam.	Page
Figures 1, 2.—	<i>Trochammina pauciloculata</i> , Brady,	× 50	344
	<i>a, a</i> , Superior lateral aspect; <i>b, b</i> , inferior lateral aspect; <i>c, c</i> , peripheral aspect.		
,, 3.—	<i>Trochammina squamata</i> , Jones and Parker,	× 50	337
	<i>a</i> , Superior aspect; <i>b</i> , inferior aspect; <i>c</i> , peripheral aspect.		
,, 4.—	<i>Trochammina inflata</i> , Montagu, sp.,	× 50	338
	<i>a</i> , Superior aspect; <i>b</i> , inferior aspect; <i>c</i> , peripheral aspect.		
,, 5, 6.—	<i>Trochammina nitida</i> , Brady,	× 50	339
	Figs. 5 <i>a, 6</i> . Superior aspect; <i>b</i> , inferior aspect; <i>c</i> , peripheral aspect.		
,, 7-10.—	<i>Carterina spiculotesta</i> , Carter, sp.,		346
	Fig. 7 <i>a</i> , Superior aspect; <i>b</i> , inferior aspect,	× 50	
	Fig. 8. Young specimen,	× 50	
	Figs. 9, 10. Fragments of the test mounted in Canada balsam, showing the form and arrangement of the spicules,	× 100	
,, 11.—	<i>Webbina hemisphaerica</i> , Jones, Parker, and Brady,	× 25	350
,, 12-16.—	<i>Webbina clavata</i> , Jones and Parker,	× 25	349
	Figs. 12, 13. Adherent specimens.		
	Fig. 14. Small specimen with orifice on the superior surface.		
	Fig. 15. Inferior aspect, showing the cavity of the test.		
	Fig. 16. Long tubular neck, adherent to <i>Globigerinae</i> .		