

PLATE XLVII.

VERNEUILINA.

	Diam.	Page
Figures 1-3.— <i>Verneuilina spinulosa</i> , Reuss,	× 70	384
Figs. 1 <i>a</i> , 2, 3. Lateral aspect; 1 <i>b</i> , oral aspect.		
„ 4-7.— <i>Verneuilina pygmæa</i> , Egger, sp.,		385
Figs. 4, 6, Lateral aspect,		
Fig. 5. <i>a</i> , Lateral aspect; <i>b</i> , oral aspect,		
Fig. 7. Section of the shell, showing its finely arenaceous structure,		
	× 35	
	× 25	
	× 120	
„ 8-14.— <i>Verneuilina propinqua</i> , n. sp.,		387
Figs. 8, 9. Lateral aspect,		
Fig. 10. Lateral aspect; fig. 11, oral aspect,		
Fig. 12. Transverse section of the test.		
Figs. 13, 14. Adherent specimens,		
	× 20	
	× 14	
	× 20	
„ 15-17.— <i>Verneuilina polystropha</i> , Reuss, sp.,	× 60	386
Figs. 15, 16. Lateral aspect; fig. 17, oral aspect.		
„ 18-20.— <i>Verneuilina triquetra</i> , Münster, sp.,		383
Fig. 18. <i>a</i> , Lateral aspect; <i>b</i> , oral aspect,		
Figs. 19, 20 <i>a</i> . Lateral aspect; <i>b</i> , oral aspect,		
	× 14	
	× 20	
„ 21-24.— <i>Verneuilina variabilis</i> , n. sp.,	× 14	385
<i>a, a, a</i> , Lateral aspect of specimens of various forms;		
<i>b, b, b</i> , oral aspect of the same.		