

PLATE LXXXII.

GLOBIGERINA.—ORBULINA.—CANDEINA.

a, Superior aspect; *b*, inferior aspect; *c*, peripheral aspect.

	Diam.	Page
Figures 1-3.— <i>Orbulina universa</i> , d'Orbigny,	× 50	608
Fig. 1. Spinous pelagic specimen.		
Fig. 2. Two-chambered shell laid open.		
Fig. 3. Specimen with two supplementary chambers.		
„ 4.— <i>Globigerina sacculifera</i> , Brady,	× 50	604
Adult pelagic specimen.		
„ 5.— <i>Globigerina conglobata</i> , Brady,	× 50	603
Adult pelagic specimen.		
„ 6, 7.— <i>Globigerina digitata</i> , Brady,	× 50	599
Large bottom specimens.		
„ 8, 9.— <i>Globigerina</i> , sp.,	× 50	603
Minute <i>Globigerinæ</i> ; general contour resembling that of <i>Globigerina elevata</i> , d'Orbigny, and <i>Globigerina trochoides</i> , Reuss.		
„ 10.— <i>Globigerina (cretacea)</i> , d'Orbigny?,	× 50	596
Bottom specimen.		
„ 11.— <i>Globigerina cretacea</i> , d'Orbigny,	× 50	596
Fossil specimen.		
„ 12.— <i>Globigerina linnæana</i> , d'Orbigny, sp.,	× 50	598
Fossil specimen.		
„ 13-20.— <i>Candeina nitida</i> , d'Orbigny,		622
Fig. 13. Pelagic specimen,	× 50	
Figs. 14-18. Bottom specimens; figs. 14, 18, superior aspect; fig. 15, inferior aspect; figs. 16, 17, peripheral aspect,	× 50	
Fig. 19. Monstrous specimen,	× 50	
Fig. 20. Portion of the shell more highly magnified, showing the sutural apertures and the minute perforation of the walls,	× 100	