

PLATE CIV.

PULVINULINA.

Figures		Diam.	Page
1, 2.	<p>—<i>Pulvinulina micheliniana</i>, d'Orbigny, sp., .</p> <p><i>a</i>, Superior aspect; <i>b, b</i>, inferior aspect; <i>c, c</i>, peripheral aspect.</p> <p>Fig. 1. Bottom specimen; fig. 2, surface specimen.</p>	× 40	694
,,	<p>3-11.—<i>Pulvinulina pauperata</i>, Parker and Jones, .</p> <p>Figs. 3-9. Lateral aspects of the shell, in various stages.</p> <p>Figs. 10-11. Peripheral aspect.</p>	× 25	696
,,	<p>12-16.—<i>Pulvinulina favus</i>, Brady, . . .</p> <p>Figs. 12 <i>a</i>, 13 <i>a</i>, Lateral aspect; figs. 12 <i>b</i>, 13 <i>b</i>, peripheral aspect,</p> <p>Fig. 14. Horizontal section of the test,</p> <p>Figs. 15, 16. Portions of the surface more highly magnified,</p>	<p>× 40</p> <p>× 40</p> <p>× 80</p>	701
,,	<p>17.—<i>Pulvinulina punctulata</i>, d'Orbigny, sp., .</p> <p><i>a</i>, Superior aspect; <i>b</i>, inferior aspect; <i>c</i>, peripheral aspect.</p>	× 16	685
,,	<p>18.—<i>Pulvinulina repanda</i>, Fichtel and Moll, sp., .</p> <p><i>a</i>, Superior aspect; <i>b</i>, inferior aspect; <i>c</i>, peripheral aspect.</p>	× 30	684
,,	<p>19.—<i>Pulvinulina repanda</i>, var. <i>concamerata</i>, Montagu, var.,</p> <p><i>a</i>, Superior aspect; <i>b</i>, inferior aspect; <i>c</i>, peripheral aspect.</p>	× 30	685