

PLATE CVII.

ROTALIA.

Figure	Diam.	Page
1.— <i>Rotalia papillosa</i> , var. <i>compressiuscula</i> , nov.,	× 30	708
<i>a</i> , Superior aspect; <i>b</i> , inferior aspect; <i>c</i> , peripheral aspect.		
2, 3.— <i>Rotalia beccarii</i> , Linné, sp.,	× 50	704
<i>a,a</i> , Superior aspect, <i>b,b</i> , inferior aspect; <i>c,c</i> , peripheral aspect.		
4.— <i>Rotalia broeckhiana</i> , Karrer,	× 50	705
<i>a</i> , Superior aspect; <i>b</i> , inferior aspect; <i>c</i> , peripheral aspect.		
5.— <i>Rotalia orbicularis</i> (?), d'Orbigny,	× 50	706
<i>a</i> , Superior aspect; <i>b</i> , inferior aspect; <i>c</i> , peripheral aspect. (A passage-form between <i>Rotalia beccarii</i> and <i>Rotalia orbicularis</i> . The typical <i>Rotalia orbicularis</i> is given at Pl. CXV. fig. 6.)		
6, 7.— <i>Rotalia soldanii</i> , d'Orbigny,	× 40	706
<i>a,a</i> , Superior aspect; <i>b,b</i> , inferior aspect; <i>c,c</i> , peripheral aspect.		
8, 9.— <i>Rotalia clathrata</i> , n. sp.,	× 50	709
<i>a,a</i> , Superior aspect; <i>b,b</i> , inferior aspect; <i>c,c</i> , peripheral aspect.		