

PLATE CXV.

PULVINULINA.—ROTALIA, &c.

		Diam.	Page
Figure 1.—	<i>Pulvinulina schreibersii</i> , d'Orbigny, n. sp.,	× 30	697
	<i>a</i> , Superior aspect ; <i>b</i> , inferior aspect ; <i>c</i> , peripheral aspect.		
„ 2.—	<i>Pulvinulina vermiculata</i> , d'Orbigny, sp.,	× 30	687
	<i>a</i> , Superior aspect ; <i>b</i> , inferior aspect.		
„ 3.—	<i>Pulvinulina dispansa</i> , n. sp.,	× 14	687
	<i>a</i> , Superior aspect ; <i>b</i> , inferior aspect.		
„ 4, 5.—	<i>Truncatulina lobatula</i> , Walker and Jacob, sp.,	× 30	660
	Fig. 4. Adherent specimen enclosed in a nidus of sand. Fig. 5. Similar specimen, the sand partially removed.		
„ 6.—	<i>Rotalia orbicularis</i> , d'Orbigny,	× 50	706
	<i>a</i> , Superior aspect ; <i>b</i> , inferior aspect ; <i>c</i> , peripheral aspect.		
„ 7.—	<i>Rotalia schroeteriana</i> , Parker and Jones,	× 25	707
	<i>a</i> , Superior aspect ; <i>b</i> , inferior aspect ; <i>c</i> , peripheral aspect.		
„ 8.—	<i>Rotalia pulchella</i> , d'Orbigny, sp.,	× 50	710
	<i>a</i> , Superior aspect ; <i>b</i> , inferior aspect.		
„ 9.—	<i>Nonionina boueana</i> , var. <i>armata</i> , nov.,	× 60	730
	<i>a</i> , Lateral aspect ; <i>b</i> , peripheral aspect.		