

The following is a list of the species :—

1. *Ceratosoma cornigerum* (Adams), Bergh.
Indian Ocean, Philippine Sea.
2. *Ceratosoma gracillimum*, Semper.
Philippine Sea.
3. *Ceratosoma trilobatum*, Gray.
Red Sea.
4. *Ceratosoma polyomma*, Bergh.
Pacific Ocean (Pelew Islands).
5. *Ceratosoma caledonicum*, Fischer.
Pacific (New Caledonia).
6. *Ceratosoma brevicaudatum*, Abraham.
Pacific.
7. *Ceratosoma oblongum*, Abraham.
Pacific (West Australia).
8. *Ceratosoma tenue*, Abraham.
Habitat ?

Ceratosoma cornigerum (Adams), Bergh, var. ? (Pl. II. figs. 14–17; Pl. III. figs. 14–20).

Ceratosoma cornigerum, A. Adams, Voyage of the "Samarang," Mollusca, 1848, p. 68, pl. xix. fig. 5.
 „ Adams, Bergh, Malacolog. Untersuch, *loc. cit.*, Heft x., 1876, pp. 393–403,
 Taf. xlvi. figs. 15–27, Taf. xlix. figs. 1–5.

Habitat.—Philippine Sea (Samboangan).

One specimen of this species was dredged at Samboangan, in company with *Platydoris eurychlamys* and *Discodoris morphæa*, on February 1, 1875, from a depth of 10 fathoms. The specimen was well preserved in alcohol, and measured 4·3 cm. in length by 1 in breadth and 1·5 cm. in height; the hinder dorsal process, moreover, measured 6 mm.; the breadth of the foot 5 mm., the length of the tail 16 mm., the height of the rhinophoria 3 mm., of the branchia 3·5 mm. The colour was greyish-red, white speckled and covered with numerous white dots, especially on the anterior portion of the back. The stem of the rhinophoria