

Species of <i>Myzostoma</i> .	Host.	Received from.
<i>M. tenuispinum</i> , n. sp.	{ <i>Antedon inæqualis</i> , P. H. C.	Challenger Expedition.
	„ <i>incisa</i> , P. H. C.	Do. do.
	„ <i>angusticalyx</i> , P. H. C.	Do. do.
	„ <i>basicurva</i> , P. H. C.	Do. do.
<i>M. willemoesii</i> , n. sp.	{ <i>Antedon basicurva</i> , P. H. C.	Do. do.
	„ <i>inæqualis</i> , P. H. C.	Do. do.
<i>M. inflator</i> , n. sp.	{ <i>Antedon angustiradia</i> , P. H. C.	Do. do.
	{ <i>Actinomeiru pulchella</i> , Pourt., sp.	“Blake.”
<i>M. murrayi</i> , n. sp.	{ <i>Antedon angustiradia</i> , P. H. C.	Challenger Expedition.
	„ <i>radiospina</i> , P. H. C.	Do. do.
	„ <i>duplex</i> , P. H. C.	“Blake.”
<i>Stelechopus hyocrini</i> , n. g., n. sp.	<i>Hyocrinus</i> , sp. ?	Challenger Expedition.

TABLE II.

List of Crinoids on which Myzostomida have been hitherto found.

Host.	Species of <i>Myzostoma</i> .	Living.
<i>Antedon angusticalyx</i> , P. H. C.	<i>M. tenuispinum</i> , n. sp.	In roundish arm cysts.
„ <i>angustiradia</i> , P. H. C.	{ <i>M. inflator</i> , n. sp.	In independent sessile cysts.
	{ <i>M. murrayi</i> , n. sp.	In independent stalked cysts.
„ <i>basicurva</i> , P. H. C.	{ <i>M. willemoesii</i> , n. sp.	In spiral malformations of the pinnules.
	{ <i>M. tenuispinum</i> , n. sp.	In roundish arm cysts.
„ <i>bidentata</i> , P. H. C.	{ <i>M. dentatum</i> , n. sp.	Free.
	{ <i>M. quadrifilum</i> , n. sp.	Do.
	{ <i>M. filiferum</i> , n. sp.	Do.
„ <i>carinata</i> , Leach, sp.	<i>M. gigas</i> , Lütken.	Do.