

PLATE XVII.¹

SCHIZOPORELLA.—SMITTIA.—LEPRALIA.—CRIBRILINA.—MUCRONELLA.

	PAGE
Figure 1.— <i>Schizoporella nivea</i> ,	163
1 <i>a</i> , magnified.	
„ 2.— <i>Schizoporella longispinata</i> ,	163
2 <i>a</i> , magnified; 2 <i>b</i> , single zoecium, more highly magnified; 2 <i>c</i> , oecium.	
„ 3.— <i>Smittia smittiana</i> ,	151
3 <i>a</i> , magnified; 3 <i>b</i> , oecium.	
„ 4.— <i>Lepralia celleporoides</i> ,	142
4 <i>a</i> , magnified; 4 <i>b</i> , more highly magnified.	
„ 5.— <i>Lepralia japonica</i> ,	143
5 <i>a</i> , magnified.	
„ 6.— <i>Cribrilina philomela</i> ,	132
6 <i>a</i> , magnified; 6 <i>b</i> , oecium.	
„ 7.— <i>Lepralia tuberosa</i> ,	143
7 <i>a</i> , magnified; 7 <i>b</i> , with oecium; 7 <i>c</i> , single zoecium, more highly magnified; 7 <i>d</i> , back view.	
„ 8.— <i>Mucronella quadrata</i> ² (Pl. XVIII. fig. 5),	156
8 <i>a</i> , magnified.	

¹ When the plate was prepared all these forms were included under the generic term *Hemeschara*, which has since been discarded in accordance with more modern views.

² Altogether misnamed on the plate.