

STATION 210. January 25, 1875; off the Panglao and Siquijor Islands; lat. $9^{\circ} 26' N.$, long. $123^{\circ} 45' E.$; 375 fathoms; blue mud; bottom temperature, $54^{\circ} 1 F.$

Pentacrinus alternicirrus (?). Also at Stations 171 and 214.

Pentacrinus naresianus (?). Also at Stations 170A, 171, 175, and 214.

Metacrinus murrayi (?).

Antedon (two species).

STATION 214. February 10, 1875; off the Meangis Islands; lat. $4^{\circ} 33' N.$, long. $127^{\circ} 6' E.$; 500 fathoms; blue mud; bottom temperature, $41^{\circ} 8 F.$

Pentacrinus alternicirrus. Also at Stations 171 and 210 (?).

Pentacrinus naresianus. Also at Stations 170A, 171, 175, and 210 (?).

Metacrinus costatus.

Metacrinus moseleyi.

Metacrinus murrayi (?).

Metacrinus varians.

Metacrinus wyvillii. Also at Station 170A.

Antedon (six species).

Promachocrinus naresi.

STATION 223. March 19, 1875; lat. $5^{\circ} 31' N.$, long. $145^{\circ} 13' E.$; 2325 fathoms; Globigerina ooze; bottom temperature, $35^{\circ} 5 F.$

Hyocrinus bethellianus, young (*fide* C. W. T.). Also at Stations 106 (C. W. T.) and 147.

STATION 235. June 4, 1875; lat. $34^{\circ} 7' N.$, long. $138^{\circ} 0' E.$; 565 fathoms; green mud; bottom temperature, $38^{\circ} 1 F.$

Pentacrinus (?) *mollis*.

Eudiocrinus japonicus.

STATION 323. February 28, 1876; lat. $35^{\circ} 39' S.$, long. $50^{\circ} 47' W.$; 1900 fathoms; blue mud; bottom temperature, $33^{\circ} 1 F.$

Rhizocrinus lofotensis (*fide* C. W. T.). Also the "Lightning" (1868), Stations 12 and 16; the "Knight Errant" (1880), Stations 5 and 6; and the Challenger, Stations 24A and 122c. Also at several Stations in the Gulf Stream, the Caribbean Sea, and off the New England coast.