

STATION LIST OF THE STALKED CRINOIDS WHICH HAVE BEEN OBTAINED BY THE VARIOUS AMERICAN EXPEDITIONS FOR DEEP-SEA EXPLORATION (MOSTLY UNDER THE DIRECTION OF MR. ALEXANDER AGASSIZ) BETWEEN THE YEARS 1867 AND 1880.

The presence of Comatulæ is noticed as in the previous list ; but the occurrence of a species at several Stations is not, as a rule, recorded more than once.

SS. "CORWIN," 1867.

No. 2. May 24, 1867 ; 1.6 miles from Chorrera, Cuba ; 270 fathoms.

*Pentacrinus decorus* (stem only ?).

SS. "BIBB," 1868-69.

May 9, 1868 ; off the Samboes ; 237 fathoms.

May 11, 1868 ; off Sand Key ; 248 and 306 fathoms.

*Rhizocrinus lofotensis*. Also off Havana, Barbados, Grenada, and Grenadines. Dredged by the "Lightning," the Challenger, and the "Knight Errant."

March 4, 1869 ; off Cojima, near Havana ; 450 fathoms.

*Rhizocrinus lofotensis*.

*Atelecrinus cubensis*.

*Antedon*.

March 10, 1869 ; off Double-headed Shot Keys ; 315 and 471 (?) fathoms.

*Pentacrinus decorus* (stem only ?).

SS. "HASSLER," 1871-72.

December 30, 1871. Off Sandy Bay, Barbados ; 100 fathoms.

*Rhizocrinus rawsoni*. Also off Havana, Montserrat, Guadeloupe, Dominica, Martinique, and Panama. Dredged in the East Atlantic by the "Porcupine," Challenger, "Talisman," and "Travailleur."

SS. "BLAKE," 1878-80.

Cruise of 1877-78.

STATION 21. Off Bahia Honda, Cuba ; lat. 23° 2' N., long. 83° 13' W. ; 287 fathoms.

*Pentacrinus decorus*. Also off Havana, Montserrat, St. Vincent Barbados, Kingston, and Saba Island.