

PLATE IIIA.

Fig. 1. Simple slender dorsal bristle, minutely roughened toward the tip, of <i>Hermodice carunculata</i> ,	x 350.
Fig. 2. One of the stouter serrated dorsal bristles of the same,	x 350.
Fig. 3. One of the upper series of ventral bristles of the foregoing, with a more elongated tip,	x 350.
Fig. 4. A stouter ventral bristle of the same species,	x 350.
Fig. 5. Simple, slightly curved bristle of <i>Eurythoë pacifica</i> ,	x 350.
Fig. 6. One of the elongated slightly bifid dorsal bristles of the same,	x 350.
Fig. 7. Third kind (stout serrated) of dorsal bristle of the foregoing,	x 350.
Fig. 8. A long (superior) bifid ventral bristle of the same, with a finely serrated inner margin,	x 350.
Fig. 9. One of the stouter ventral bristles of the preceding, having only three or four serrations near the tip,	x 350.
Fig. 10. Elongated translucent bifid bristle of <i>Eurythoë pacifica</i> , var. <i>levukaensis</i> ,	x 350.
Fig. 11. Stout serrated dorsal bristle of the same,	x 350.
Fig. 12. Ventral bristle (of medium size) from the anterior third of the fore- going species,	x 350.
Fig. 13. Simple translucent dorsal bristle of <i>Hipponeë gaudichaudi</i> ,	x 350.
Fig. 14. Bifid tip of one of the foregoing,	x 700.
Fig. 15. Normal appearance of the lower part of the shaft of the same kind of bristle, only a slight roughness along the edge indicating the points,	x 700.
Fig. 16. A compressed (and fissured) shaft of the same form, with more distinct indications of the points or serratures,	x 700.
Fig. 17. Bifid ventral hook of the preceding species,	x 350.