

PLATE VA.

- Fig. 1. Tip of one of the great dorsal spines of *Lætmonice producta*, var. *benthaliana*, × 90.
- Fig. 2. Tip of a ventral bristle from the same form, × 55.
- Fig. 3. Tip of a short (developing?) spine from the posterior third of *Lætmonice producta*, var. *willemoesii*, × 90.
- Fig. 4. Ventral bristle of the same species, × 55.
- Fig. 5. Tip of a dorsal spine of *Lætmonice producta*, var. *assimilis*, × 90.
- Fig. 6. A more extensive view of the tip of one of the spines, so as to demonstrate the terminal curves, × 55.
- Fig. 7. A similar view of the organ in *Lætmonice flicornis*, from Shetland. The contrast in the curves is evident, × 55.
- Fig. 8. Tip of a ventral bristle of *Lætmonice producta*, var. *assimilis*, × 90.
- Fig. 9. Tip of a ventral bristle of *Lætmonice japonica*, × 55.
- Fig. 10. Tip of a dorsal spine of the same species, × 90.
- Fig. 11. Extremity of one of the very fine hairs forming the "felt" of *Lætmonice aphroditoides*, × 350.
- Fig. 12. Tip of one of the lustrous brown dorsal spines of this form, × 55.
- Fig. 13. Ventral bristle of the same species, showing the "hairy" condition of the tip, × 55.
- Fig. 14. Tip of one of the minute serrate bristles from the anterior feet of the same species, × 350.
- Fig. 15. Extremity of one of the stout brownish spines from the same region of the body, × 90.