

PLATE VIIA.

Fig. 1.	Average dorsal bristle of <i>Lagisca magellanica</i> ,	×	90.
Fig. 2.	Ventral bristle of the same (from the middle of the series),	×	90.
Fig. 3.	Tip of average dorsal bristle of <i>Eulagisca corrientis</i> ,	×	90.
Fig. 4.	Bristle from the middle of the ventral tuft of the same form,	×	90.
Fig. 5.	Average dorsal bristle of <i>Lagisca tenuisetis</i> ,	×	400.
Fig. 6.	Tip of one of the large forms from the same tuft,	×	400.
Figs. 7, 8.	Superior ventral bristles of the same species,	×	400.
Fig. 9.	Bristle with shorter tip, from the middle of the ventral series of the same,	×	400.
Fig. 10.	Tip of a dorsal bristle of <i>Lagisca peracuta</i> ,	×	350.
Fig. 11.	Extremity of one of the middle bristles of the ventral series of the same species,	×	350.
Fig. 12.	Tip of a dorsal bristle of <i>Polyeunoa lævis</i> ,	×	210.
Fig. 13.	Ventral bristle of the foregoing,	×	210.
Fig. 14.	Tip of a dorsal bristle of <i>Lagisca magellanica</i> , var. <i>grubei</i> ,	×	210.
Fig. 15.	One of the superior ventral bristles of the same form,	×	90.
Fig. 16.	Tip of a bristle from the middle of the ventral series of the foregoing,	×	210.