

PLATE XVI.

Fig. 1. Setigerous region of a foot of <i>Syllis ramosa</i> (parent stock),	x 400.
Fig. 2. Dorsal bristle, from the tenth foot of <i>Nereis (Platynereis) kobiensis</i> ,	x 400.
Fig. 3. Amber-coloured hook from the inferior border of the group, from the thirty-seventh foot of the same,	x 400.
Fig. 4. One of the larger superior falcate bristles, from the inferior division of the foregoing foot,	x 400.
Fig. 5. One of the dorsal bristles, from the tenth foot of <i>Nereis tongatabuensis</i> ,	x 400.
Fig. 6. Falcate bristle, from the superior lobe of the thirty-seventh foot of the foregoing,	x 590.
Fig. 7. Falcate bristle, from the inferior lobe of the same foot,	x 590.
Fig. 8. Dorsal bristle, from the thirty-seventh foot of <i>Nereis melanocephala</i> ,	x 400.
Fig. 9. Distal end of a falcate bristle, from the ventral series of the same foot,	x 400.
Fig. 10. Setose dorsal bristle, from the thirty-seventh foot of <i>Nereis atlantica</i> ,	x 400.
Fig. 11. A falcate bristle, from the inferior division of the tenth foot of the same,	x 400.
Fig. 12. Setose dorsal bristle, from the tenth foot of <i>Nereis longisetis</i> ,	x 400.
Fig. 13. One of the inferior ventral bristles, with very short setose tips, in the twenty-seventh foot of the foregoing,	x 400.
Fig. 14. A setose bristle, from the dorsal series of <i>Nereis antillensis</i> ,	x 400.
Fig. 15. A falcate bristle, from the superior division of one of the posterior feet of the same,	x 400.
Fig. 16. One of the upper falcate forms, from the ventral division of the foot posteriorly,	x 400.
Fig. 17. Setose dorsal bristle, from the tenth foot of <i>Nereis kerguelensis</i> ,	x 400.
Fig. 18. One of the upper falcate bristles, from the inferior series of the tenth foot of the same,	x 530.
Fig. 19. One of the falcate bristles, from the inferior division of the foot of <i>Nereis longisetis</i> ,	x 400.