

PLATE XVIII.

- Fig. 1. A lateral view of one of the posterior hooks of *Lumbriconereis japonica*, × 250.
- Fig. 2. One of the winged bristles next the winged hooks in the tenth foot of *Lumbriconereis kerguelensis*, × 250.
- Fig. 3. The upper hook of the series in the tenth foot of the same, × 400.
- Fig. 4. Two of the extremely elongated bristles projecting from the thirtieth foot of the foregoing, × 90.
- Fig. 5. One of the elongated bristles, from the sixteenth foot of *Lumbriconereis neo-zealandia*, var. A, × 250.
- Fig. 6. Posterior hook of the same, var. A, × 400.
- Fig. 7. Posterior (and superior) hook of *Lumbriconereis neo-zealandia*, var. B, × 400.
- Fig. 8. Winged bristle, from the sixteenth segment of the same (var. B), × 250.
- Fig. 9. The third winged hook (viewed antero-posteriorly) of the tenth foot of the foregoing (var. B), × 250.
- Fig. 10. Posterior hook of *Lumbriconereis abyssorum*, × 400.
- Fig. 11. Posterior hook of *Lumbriconereis punctata*, × 400.
- Fig. 12. Posterior hook of *Lumbriconereis ehlersi*, var. *tenuisetis*, × 400.
- Fig. 13. An average hook, from the fragmentary example of *Lumbriconereis heteropoda*, × 250.
- Fig. 14. Tip of a bristle (from the group below the dorsal hook) of the foregoing, × 250.
- Fig. 15. A posterior bristle of *Notocirrus capensis*, × 250.
- Fig. 16. Dorsal bristle, from the thirtieth foot of *Nematonereis schmarda*, × 400.
- Fig. 17. Jointed bristle, from the thirtieth foot of the same, × 400.
- Fig. 18. Upper brownish spine (with hook and wing), from the fiftieth foot of the foregoing, × 400.
- Fig. 19. Brush-like bristle, from the posterior region of the same, × 400.
- Fig. 20. One of the stouter bristles, from a fragment of a male specimen of *Palolo viridis*, × 400.
- Fig. 21. A jointed bristle from the same specimen, × 400.