

STATION 208.—January 17, 1875; lat. $11^{\circ} 37' N.$, long. $123^{\circ} 31' E.$; depth, 18 fathoms; bottom, blue mud (same as Station 204).

Chama carditæformis, Reeve.
Anatina siphonata, Reeve.
Cardium (Bucardium) tenuicos-
tatum, Lamarck.
Cardium (Bucardium) australe,
 Sowerby.

Diplodonta subgranulosa, n. sp.
Arca (Barbatia) lima, Reeve.
Pecten leopardus (var. *solaris*).
Pecten senatorius, var.

STATION 209.—January 22, 1875; lat. $10^{\circ} 14' N.$, long. $123^{\circ} 54' E.$; depth, 95 fathoms; bottom, blue mud (east of the Island of Panay, Philippines).

Pecten vitreus, Chemnitz.

STATION 210.—January 25, 1875; lat. $9^{\circ} 26' N.$, long. $123^{\circ} 45' E.$; depth, 375 fathoms; bottom, blue mud (same position as Station 209).

Semele (Abra) philippinensis, n. sp. | *Amussium jeffreysi*, n. sp.

STATION 212.—January 30, 1875; lat. $6^{\circ} 54' N.$, long. $122^{\circ} 18' E.$; depth, 10 fathoms; bottom, sand (south of the Island of Mindanao).

Cytherea (Callista) lilacina,
 Lamarck.
Cytherea (Callista) roseotincta, n. sp.
Cytherea (Caryatis) hebræa,
 Lamarck.
Circe sulcata, Gray.
Venus (Chione) recognita, n. sp.

Mactra (Trigonella) incarnata,
 Deshayes.
Cardium setosum, Redfield.
Cardita canaliculata, Reeve.
Arca (Scapharca) angicostata,
 Reeve.
Pectunculus striatularis, Lamarck (?).

Lima squamosa, Lamarck.

STATION 216.—February 16, 1875; lat. $2^{\circ} 46' N.$, long. $133^{\circ} 58' E.$; depth, 1675 fathoms; bottom, Globigerina ooze (north-west of New Guinea).

Arca (Barbatia) corpulenta, n. sp.

STATION 218.—March 1, 1875; lat. $2^{\circ} 23' S.$, long. $144^{\circ} 4' E.$; depth, 1070 fathoms; bottom, blue mud (north of New Guinea).

Amussium watsoni, n. sp.

STATION 219.—March 10, 1875; lat. $1^{\circ} 54' S.$, long. $146^{\circ} 39' 40'' E.$; depth, 150 fathoms; bottom, coral mud (a little north-east of Station 218).

Cryptodon watsoni, n. sp.