

SECTION B.—Subgenus *Cardiomya*, A. Adams = *Spathophora*, Jeffreys.

Teeth.—Same as in (*Neæra* proper) Section A.

Cartilage.—Less oblique than in the above.

Surface.—Radiately costate or lirate.

Neæra gouldiana, Hinds.

costellata, Deshayes.

fallax, n. sp.

pectinata, Carpenter.

costata, Sowerby.

Neæra concinna, Hinds.

striata, Jeffreys.

curta, Jeffreys.

multicostata, Verrill and
Smith.*

SECTION C.

Teeth of right valve, *Cartilage-Pit*, and *Sculpture*, the same as in Section A, but a small erect posterior lateral tooth is present in the left valve.

Neæra pulchella, H. Adams.

SECTION D.

Teeth.—One small lateral tooth on each side near the beak in both valves (the anterior in the left being feeble).

Cartilage and *Surface* as in A and B.

Neæra singaporensis, Hinds.

SECTION E.

Teeth.—A short lateral tooth on each side in the right valve, with corresponding pits in the hinge-margin of the left, more conspicuous on the anterior side than behind.

Cartilage.—Obliquely inclined towards the posterior end.

Surface.—Concentrically striated.

Neæra teres, Jeffreys.

SECTION F.

Teeth.—A short erect lateral tooth on each side, and a small dentiform projection of the hinge-margin in front of the apex in the right valve, and a single tubercular denticle in front of the apex (but no laterals) in the left.