

TABLE SHOWING THE DISTRIBUTION OF THE FAMILIES, GENERA, AND SPECIES OF ASCIDIÆ COMPOSITÆ<sup>1</sup> THROUGHOUT THE GREAT OCEAN BASINS.

SPECIES.	1. North Atlantic.	2. South Atlantic.	3. Southern Ocean.	4. Seas of Malay Archip.	5. North Pacific.	6. South Pacific.	7. Seas of South America.
Family BOTRYLLIDÆ.							
<i>Botrylloides tyreum</i> , . . . . .				+			
<i>perspicuum</i> , . . . . .				+			
var. <i>rubicundum</i> , . . . . .				+			
<i>nigrum</i> , . . . . .	+						
<i>fulgurale</i> , . . . . .	+						
<i>Sarcobotrylloides wyvillii</i> , . . . . .	+						
<i>Polycyclus lamarcki</i> , . . . . .	+						
<i>jeffreysi</i> , . . . . .	+						
Family DISTOMIDÆ.							
<i>Colella pedunculata</i> , . . . . .			+				+
<i>thomsoni</i> , . . . . .				+			
<i>gaimardi</i> , . . . . .							+
<i>pulchra</i> , . . . . .				+			
<i>elongata</i> , . . . . .						+	
<i>quoyi</i> , . . . . .			+				
<i>murrayi</i> , . . . . .						+	
var. <i>rubida</i> , . . . . .						+	
<i>ramulosa</i> , . . . . .							+
<i>concreta</i> , . . . . .			+				
<i>Distaplia vallii</i> , . . . . .	+			+			
— (f) <i>clava</i> , . . . . .	+						
— (f) <i>pyriformis</i> , . . . . .			+				
<i>Cystodytes draschii</i> , . . . . .		+					
<i>philippinensis</i> , . . . . .				+			
<i>Symplegma viride</i> , . . . . .	+						
Family POLYOLINIDÆ.							
<i>Pharyngodictyon mirabile</i> , . . . . .			+				
<i>Tylobranchion speciosum</i> , . . . . .			+				
<i>Atopogaster gigantea</i> , . . . . .							+
<i>aurantiaca</i> , . . . . .						+	
<i>informis</i> , . . . . .							
<i>elongata</i> , . . . . .							+
var. <i>pallida</i> , . . . . .		+					
<i>Morchellioides affinis</i> , . . . . .			+				
<i>Morchellium giardi</i> , . . . . .			+				

<sup>1</sup> Those collected during the "Lightning," "Porcupine," and Challenger Expeditions.