

SPECIES.	1. North Atlantic.	2. South Atlantic.	3. Southern Ocean.	4. Seas of Malay Archip.	5. North Pacific.	6. South Pacific.	7. Seas of South America.
<i>Leptoclinum speciosum</i> , .. . . .		+					
var. <i>asperum</i> , . . . . .		+	+				
<i>annectens</i> , . . . . .		+					
<i>tenuis</i> , . . . . .	+	+					+
var. <i>magnizoidium</i> , . . . . .	+						
<i>propinquum</i> , . . . . .							+
<i>neglectum</i> , .. . . .							
<i>albidum</i> (?), . . . . .	+	+					
var. <i>luteolum</i> (?), . . . . .	+						
var. <i>grande</i> , .. . . .				+			
<i>subflavum</i> , .. . . .			+				
<i>jeffreysi</i> , .. . . .	+						
<i>carpenteri</i> , . . . . .	+						
<i>thomsoni</i> , . . . . .	+						
<i>edwardsi</i> , . . . . .		+					
<i>japonicum</i> , . . . . .					+		
<i>Leptoclinum</i> (?) <i>jacksoni</i> , . . . . .						+	
<i>rubicundum</i> , . . . . .			+				
Family DIPLOSOMIDÆ.							
<i>Diplosomoides molle</i> , . . . . .				+			
<i>Diplosoma macdonaldi</i> , . . . . .		+					
Family CÆLOCORMIDÆ.							
<i>Cælocormus huxleyi</i> , . . . . .							+
Family POLYSTYELIDÆ.							
<i>Goodsiria placenta</i> , . . . . .		+					
var. <i>fusca</i> , . . . . .		+					
<i>pedunculata</i> , . . . . .							+
<i>coccinea</i> , . . . . .							+
<i>Synstyela incrustans</i> , . . . . .				+			+
<i>Chorizocormus reticulatus</i> , . . . . .			+				
Total number of species or varieties which occurred in each of the seven areas }	18	17	24	15	1	10	21