

North Latitude.					SPECIES.	South Latitude.				
65°-40°	40°-30°	30°-20°	20°-10°	10°-0°		0°-10°	10°-20°	20°-30°	30°-40°	40°-55°
+					<i>Aplidium fallax</i> , . . . .					
+				+	<i>despectum</i> , . . . .					+
					<i>fumigatum</i> , . . . .					+
					<i>Amarouciump variabile</i> , . . . .					+
					var. <i>tenerum</i> , . . . .					+
					<i>globosum</i> , . . . .					+
					<i>complanatum</i> , . . . .					+
					<i>irregulare</i> , . . . .					+
					v. <i>concinnum</i> , . . . .					+
					<i>pallidulum</i> , . . . .					+
					<i>recumbens</i> , . . . .					+
					<i>hepaticum</i> , . . . .					+
					<i>laevigatum</i> , . . . .					+
					<i>coleloides</i> , . . . .					+
					<i>albidum</i> , . . . .					+
					<i>nigrum</i> , . . . .					+
					<i>Psammoplidium spongiforme</i> , . . . .					+
					<i>effrenatum</i> , . . . .					+
					<i>rude</i> , . . . .					+
					<i>subviride</i> , . . . .					+
					<i>exiguum</i> , . . . .					+
					<i>ovatum</i> , . . . .					+
					<i>retiforme</i> , . . . .					+
					<i>flavum</i> , . . . .					+
					<i>pyriforme</i> , . . . .					+
					(?) <i>ignotus</i> , . . . .					+
					<i>Didemnum savignii</i> , . . . .					+
					<i>aurantiacum</i> , . . . .					+
					<i>inerme</i> , . . . .					+
					<i>Leptoclinum tonga</i> , . . . .					+
					<i>moseleyi</i> , . . . .					+
					<i>speciosum</i> , . . . .					+
					var. <i>asperum</i> , . . . .					+
					<i>annectens</i> , . . . .					+
					<i>tenui</i> , . . . .					+
					v. <i>magnizoidium</i> , . . . .					+
					<i>propinquum</i> , . . . .					+
					<i>neglectum</i> , . . . .					+
					<i>albidum</i> (?), . . . .					+
					var. <i>luteolum</i> , . . . .					+
					var. <i>grande</i> , . . . .					+
					<i>subflavum</i> , . . . .					+
					<i>jeffreysi</i> , . . . .					
					<i>carpenteri</i> , . . . .					
					<i>thomsoni</i> , . . . .					