

Family, Genus, and Species.	Range in Depth.
<i>Psammaplidium subviride</i> ,	150 fathoms.
<i>exiguum</i> ,	98 "
<i>ovatum</i> ,	3-11 "
<i>retiforme</i> ,	50-120 "
<i>flavum</i> ,	600 "
<i>pyriforme</i> ,	8 "
<i>(?) ignotus</i> ,	55 "
DIDEMNIDÆ—	
<i>Didemnum savignii</i> ,	150 "
<i>aurantiacum</i> ,	38 "
<i>inerme</i> ,	Shallow water.
<i>Leptoclinum tonga</i> ,	18 fathoms.
<i>moseleyi</i> ,	10 "
<i>speciosum</i> ,	7-20 "
var. <i>asperum</i> ,	7-20 "
<i>annectens</i> ,	Shallow water.
<i>tenue</i> ,	175-600 fathoms.
var. <i>magnirooidium</i> ,
<i>propinquum</i> ,	245 "
<i>neglectum</i> ,
<i>albidum</i> (?),	10-125 "
var. <i>luteolum</i> (?),	35 "
var. <i>grande</i> ,	95 "
<i>subflavum</i> ,	28 "
<i>jeffreysi</i> ,	35 "
<i>carpenteri</i> ,
<i>thomsoni</i> ,
<i>edwardsi</i> ,	150 "
<i>japonicum</i> ,	50 "
<i>(?) jacksoni</i> ,	6-15 "
<i>rubicundum</i> ,	20-60 "
DIPLOSOMIDÆ—	
<i>Diplosomoides molle</i> ,	Shallow water.
<i>Diplosoma macdonaldi</i> ,	Shallow water.
CÆLOCORMIDÆ—	
<i>Cœlocormus huxleyi</i> ,	600 fathoms.
POLYSTYELIDÆ—	
<i>Goodsiria placenta</i> ,	10-20 "
var. <i>fusca</i> ,	10-20 "
<i>pedunculata</i> ,	5-10 "
<i>coccinea</i> ,	5-70 "
<i>Synstyela incrustans</i> ,	10-55 "
<i>Chorizocormus reticulatus</i> ,	30 "