

TABLE SHOWING THE BATHYMETRICAL DISTRIBUTION OF THE HOLOTHURIOIDEA.

	Locality.	Depth in fathoms.
<i>Myriotrochus rinkii</i> , Steenstrup.	Skager Rack south of Arendal, North Atlantic Ocean and Arctic Sea.	shore to 500
<i>Echinocucumis typica</i> , Sars.	North Atlantic Ocean.	40 to 530
<i>Thyone raphanus</i> , Düb. and Kor.	North Atlantic Ocean and Mediterranean Sea.	20 to 530
<i>Stichopus challengerii</i> , n. sp.	Antarctic Ocean.	550
<i>Pælopatides appendiculatus</i> , n. sp.	Off Japan.	560
<i>Stichopus tizardi</i> , Théel.	Færøe Channel.	530 to 570
<i>Trochostoma thomsonii</i> , Dan. and Kor.	North Atlantic Ocean and Arctic Sea.	136 to 658
<i>Trochostoma</i> (?).	South Atlantic Ocean.	40 to 600
<i>Psolus murrayi</i> , n. sp.	South Atlantic Ocean.	600
<i>Psolus operculatus</i> , Pourtalès.	North and South Atlantic Oceans.	120 to 600
<i>Psolus antarcticus</i> , Philippi.	Antarctic Ocean and South Atlantic Ocean.	20 to 600
<i>Holothuria intestinalis</i> , Ascan and Rathke.	North Atlantic Ocean.	10 to 650
<i>Holothuria tremula</i> , Gunner.	North Atlantic Ocean.	20 to 672
<i>Ankyroderma marenzelleri</i> , n. sp.	Off New Zealand.	700
<i>Trochostoma albicans</i> , var. <i>glabra</i> , n. sp.	Off New Zealand.	700
<i>Pælopatides aspera</i> , n. sp.	Off Philippine Islands.	700
<i>Trochostoma violaceu</i> , Studer.	Kerguelen and off New Zealand.	20 to 700
<i>Holothuria lactea</i> , n. sp.	Atlantic Ocean and off New Zealand.	700 to 1000
<i>Thyonidium pellucidum</i> , Flemming.	North Atlantic Ocean.	30 to 1081
<i>Holothuria murrayi</i> (var. ?).	North Atlantic Ocean.	1090
<i>Acanthotrochus mirabilis</i> , Dan. and Kor.	Between Norway and Spitzbergen.	658 to 1110
<i>Trochostoma albicans</i> , n. sp.	South-east of New York.	1240
<i>Stichopus torvus</i> , n. sp.	South Pacific Ocean.	1375
<i>Holothuria murrayi</i> , n. sp.	South Pacific Ocean.	1375
<i>Cucumaria abyssorum</i> , var. <i>hyalina</i> , n. sp.	Antarctic Ocean and South Pacific Ocean.	1375 to 1600
<i>Cucumaria abyssorum</i> , n. sp.	Antarctic Ocean.	1600 to 1975
<i>Pseudostichopus villosus</i> , var. <i>violaceus</i> , n. sp.	Antarctic Ocean.	1975
<i>Trochostoma</i> , sp. ?	West of Valparaiso.	2160
<i>Pælopatides confundens</i> , n. sp.	South Pacific Ocean.	1375 to 2225
<i>Cucumaria abyssorum</i> , var. <i>grandis</i> , n. sp.	South Pacific Ocean.	1500 to 2225
<i>Synapta abyssorum</i> , n. sp.	Atlantic Ocean.	2350
<i>Pseudostichopus villosus</i> , n. sp.	Antarctic Ocean, Atlantic Ocean, and Pacific Ocean.	1375 to 2900
<i>Holothuria thomsoni</i> , n. sp.	North Pacific Ocean.	1875 to 2900