

*Holothuria cavolini*, Delle Chiaje, 1823 and 1841.

*Habitat*.—Mediterranean Sea (Delle Chiaje).

*Sporadipus maculatus*, Grube, 1840.

*Habitat*.—Mediterranean Sea (Grube).

*Stichopus cinerascens*, Grube, 1840.

*Habitat*.—Mediterranean Sea (Grube).

*Holothuria lilla*, Lesson, 1830.

*Habitat*.—Borabora in Society Islands (Lesson).

*Holothuria subrubra*, Quoy and Gaimard, 1833.

*Habitat*.—Mauritius (Q. and G., Hoffmann).

*Holothuria maxima* (*Fistularia*), Forskaal, 1775 ; Selenka, 1867.

*Habitat*.—Red Sea (Forskaal, Rüppel), Mozambique (Bell).

*Holothuria obscura*, Lesueur, 1824. *Actinopyga obscura*, Verrill, 1867 to 1871.

*Habitat*.—St. Bartholomew (Lesueur).

Compare the above description of *Holothuria lubrica*.

*Holothuria aglutinata*, Lesueur, 1824. *Bohadschia agglutinata*, Verrill, 1867 to 1871.

*Habitat*.—St. Bartholomew (Lesueur).

*Holothuria nigra*, Peach, 1845 ; Foot, 1860 ; Bell, 1884. *Cucumaria niger*, Kinahan, 1859.

*Habitat*.—British Islands (Peach, Bell, Norman, Kinahan).

*Holothuria sordida* (subgenus *Microthele*), Brandt, 1835 ; Ludwig, 1881.

*Habitat*.—Caroline Islands (Brandt).

*Holothuria maculata*, Lesueur, 1824.

*Habitat*.—St. Bartholomew (Lesueur).

*Holothuria tigris* (subgenus *Microthele*), Brandt, 1835 ; Ludwig, 1881.

*Habitat*.—Caroline Islands (Brandt).