

*Holothuria* sp. (*Fistularia*), Forskaal, 1775.

*Habitat*.—Red Sea (?).

*Holothuria* sp., Anderson, 1862.

*Habitat*.—Shetland (Anderson).

Possibly a *Colochirus*.

*Holothuria fasciata*, Quoy and Gaimard, 1833.

*Habitat*.—Vanikoro (Q. and G.).

*Holothuria lucifugus*, Quoy and Gaimard, 1833. *Stichopus lucifugus*, Brandt, 1835.

*Habitat*.—Carteret (Q. and G.).

*Holothuria ophidiana*, Quoy and Gaimard, 1833.

*Habitat*.—New Guinea (Q. and G.).

*Holothuria umbrina*, Rüppell, 1828.

*Habitat*.—Red Sea (Rüppell).

*Ananus holothuroides*, Sluiter, 1880.

*Habitat*.—Java (Sluiter).

Probably a deformed individual of *Holothuria pyxis* or some other species.

*Holothuria timama*, Lesson, 1830. *Psolus timama*, Jæger, 1833.

*Habitat*.—Offack in Waigiou (Lesson).

*Holothuria forskalii*, Delle Chiaje, 1823, 1841.

*Habitat*.—Mediterranean Sea (Delle Chiaje).

*Holothuria glaberrima*, Risso, 1826.

*Holothuria ovata*, Risso, 1826.

*Holothuria mamillata*, Risso, 1826.

*Holothuria littoralis*, Risso, 1826.

*Holothuria stellata*, Risso, 1826.

*Holothuria punctata*, Risso, 1826.

These six forms were probably dredged in the Mediterranean Sea. Some of them are probably not Holothurids but belong to other groups of animals.