

TABLE I.—*continued.*

	Arctic Ocean.				Antarctic Ocean.
	North-eastern part of North America, southwards to Maine.	Greenland, Iceland, and Jan Mayen.	Northern part of Scandinavia, north of the Arctic Circle.	Rest of N. European Continent, incl. Spitzbergen, Bear Isl., Nova Zembla, and the N. Asiatic Coast to Behring Strait.	
<i>Cucumaria kerguelensis</i> , n. sp.,					x
<i>serrata</i> , n. sp.,					x
var. <i>intermedia</i> , n.,					x
var. <i>marionensis</i> , n.,					x
<i>Echinocucumis typica</i> , Sars,			x	x	
<i>Thyone spectabilis</i> , Ludwig,			x 8		x
<i>fusus</i> , O. F. Müller,					
<i>meridionalis</i> , Bell,					x
<i>muricata</i> , Studer,					x
<i>cigarro</i> , Troschel,	x				
<i>recurvata</i> , n. sp.,					x
<i>lechleri</i> , Lampert,					x
<i>Stereoderma unisemita</i> , Ayres,	x				
<i>Thyonidium drummondii</i> , Thompson,			x 8		
<i>productum</i> , Ayres,	x 8		x	x	
<i>pellucidum</i> , Fleming,			x	x	
<i>Orcula barthii</i> , Troschel,	x	x		x	
<i>luminosa</i> , Lampert,	x	x		x	
<i>Psolus regalis</i> , Verrill,	x	x	x	x	
<i>phantapus</i> , Strussenfeldt,	x	x	x	x	
<i>fabricii</i> , Düben and Koren,	x	x	x	x	
<i>squamatus</i> , Düben and Koren,	x		x		x
<i>ephippifer</i> , Wyville Thomson,					x
<i>antarcticus</i> , Philippi,					x
<i>poriferus</i> , Studer,					x
<i>disciformis</i> , n. sp.,					x
<i>incertus</i> , n. sp.,					x
<i>operculatus</i> , Pourtalès,	x 8				
<i>Stichopus sitchensis</i> , Brandt,					x
<i>challengeri</i> , n. sp.,					x
<i>Pseudostichopus mollis</i> , n. sp.,					x
<i>villosus</i> , n. sp.,					x
var. <i>violaceus</i> , n.,					x
<i>Holothuria intestinalis</i> , Ascanius and Rathke,			x	x	
<i>magellani</i> , Ludwig,					x
<i>thomsoni</i> , var. <i>hyalina</i> , n.,			x	x	
<i>ecalcarea</i> , Sars,			x ?	x ?	
<i>tremula</i> , Gunner,			x ?	x ?	