

TABLE II.—*continued.*

	Scandinavia, south of the Arctic Circle.	British Islands and France.	Spain, Portugal, Azores Island.	Mediterranean Sea.	East Coast of North America, from Maine to Florida, including Bermudas.	West Indies, incl. Florida, G. of Mexico, East Coast of Central America and South America, as far as the River Amazon.	West Coast of Africa, including Madeira, Canary Islands, &c.	East Coast of South America."
<i>Cucumaria frondosa</i> , var. <i>mediterranea</i> , Semper, <i>köllikeri</i> , Semper,				x			x	
<i>syracusana</i> , Grube,			x	x				
<i>punctata</i> , Ludwig,	x?			x?				
<i>nobilis</i> , Ludwig,	x?			x?				
<i>perepicua</i> , Ludwig,	x?			x?				
<i>minuta</i> , Fabricius,	x	x		x?	x			
<i>assimilis</i> , Düben and Koren,	x	x		x?	x			
<i>lefevrii</i> , Barrois,	x	x		x?	x			
<i>saxicola</i> , Brady and Robertson,	x	x		x?	x			
<i>cognata</i> , Lampert,	x	x		x?	x			
<i>Echinocucumis typica</i> , Sars,	x	x		x?	x			
<i>Colochirus pygmæus</i> , n. sp.,	x	x		x?	x		x	
<i>Thyone suspecta</i> , Ludwig,	x	x		x?	x		x	
<i>surinamensis</i> , Semper,	x	x		x?	x		x	
<i>briareus</i> , Lesueur, .	x	x		x?	x		x	
<i>fusus</i> , O. F. Müller,	x	x		x	x		x	
<i>scabra</i> , Verrill,	x	x		x	x		x	
<i>raphanus</i> , Düben and Koren,	x	x		x	x		x	
<i>poucheti</i> , Barrois,	x	x		x	x		x	
<i>aurantiaca</i> , Costa, .	x	x		x	x		x	
<i>gemmata</i> , Pourtalès,	x	x		x	x		x	
<i>pulcherrima</i> , Ayres,	x	x		x	x		x	
<i>peruana</i> , Lesson, .	x	x		x	x		x	
<i>glabra</i> , Ayres,	x	x		x	x		x	
<i>musculosa</i> , Ayres, .	x	x		x	x		x	
<i>inermis</i> , Heller, .	x	x		x	x		x	
<i>elongata</i> , Ayres,	x	x		x	x		x	
<i>pervicax</i> , n. sp., .	x	x		x	x		x	
<i>Stereoderma unisemita</i> , Ayres,	x	x		x	x		x	
<i>Thyonidium drummondii</i> , Thompson,	x	x		x	x		x	
<i>productum</i> , Ayres,	x	x		x	x		x	
<i>pellucidum</i> , Fleming,	x	x		x	x		x	
<i>marionii</i> , v. Marenzeller,	x	x		x	x		x	
<i>parvum</i> , Ludwig,	x	x		x	x		x	
<i>occidentale</i> , Ludwig,	x	x		x	x		x	
<i>flavum</i> , Greeff,	x	x		x	x		x	