

TABLE III.—*continued.*

	Indian Ocean.	Red Sea and Gulf of Aden.		Australia.	Pacific Ocean.		
		East Coast of Africa to Cape of Good Hope, incl. Madagascar, Mauritius, &c.	Sunda Islands, Philippine Islands, China Sea, and Bay of Bengal, Moluccas Islands, Waygeo, &c.		New Guinea, North Australia, and Queensland.	New Zealand, South Australia, New South Wales, and Victoria.	Japan Islands, Yellow Sea, and Bonin Island.
<i>Ocnus javanicus</i> , Sluiter, . . .			x				
<i>molpadiooides</i> , Semper, . . .			x				
<i>typicus</i> , n. sp., . . .			x				
<i>Echinocucumis adversaria</i> , Semper, .			x				
<i>Colochirus spinosus</i> , Quoy and Gaimard, .			x				
<i>inornatus</i> , v. Marenzeller, .		x?					
<i>quadrangularis</i> , Lesson, .		x					
<i>jagorii</i> , Semper, .		x					
<i>tristis</i> , Ludwig, .		x		x			
<i>minutus</i> , Ludwig, .			x				
<i>australis</i> , Ludwig, .			x				
<i>australis</i> , var. <i>armatus</i> , v. Marenzeller, .			x				
<i>cylindricus</i> , Semper, .			x				
<i>cucumis</i> , Semper, .			x				
<i>peruanus</i> , Semper, .			x				
<i>tuberculosus</i> , Quoy and Gaimard, .			x				
<i>viridis</i> , Semper, .			x				
<i>colloradiatus</i> , Haacke, .		x					
<i>propinquus</i> , Haacke, .		x					
<i>violaceus</i> , n. sp., .			x				
<i>challengeri</i> , n. sp., .			x				
<i>Actinocucumis typica</i> , Ludwig, .			x				
<i>difficilis</i> , Bell, .		x					
<i>Pseudocucumis acicula</i> , Semper, .	x						
<i>Amphicyclus japonicus</i> , Bell, .	x						
<i>Eucyclus duplicatus</i> , Lampert, .			x				
<i>Thyone venusta</i> , Semper, .			x				
<i>villosa</i> , Semper, .			x				
<i>buccalis</i> , Stimpson, .			x				
<i>rosacea</i> , Semper, .			x				
<i>cunninghami</i> , Bell, .			x				
<i>ovulum</i> , Selenka, .			x				
<i>mirabilis</i> , Ludwig, .			x				
<i>sacella</i> , Selenka, .	x	x	x				
<i>gibber</i> , Selenka, .			x				
<i>okeni</i> , Bell, .			x				
<i>chilensis</i> , Semper, .			x				

x

x

x

x⁸

x

x

x

x

x