

TABLE III.—*continued.*

			Indian Ocean.		Australia.		Pacific Ocean.
			East Coast of Africa to Cape of Good Hope, incl. Madagascar, Mauritius, &c.		Sunda Islands, Philippine Islands, China Sea, and Bay of Bengal, Molucca Islands, Waygeo, &c.		Japan Islands, Yellow Sea, and Bonin Island.
				Red Sea and Gulf of Aden.		New Guinea, North Australia, and Queensland.	New Zealand, South Australia, New South Wales, and Victoria.
<i>Thyone peruana</i> , Lesson,	.	.					
<i>pedata</i> , Semper,	.	.					
<i>longidentis</i> , Hutton,	.	.		x		x	
<i>brevidentis</i> , Hutton,	.	.				x	
<i>meridionalis</i> , Bell,	.	.				x	
<i>fusus</i> , var. <i>papuensis</i> , n.,	.	.	x		x		
<i>curvata</i> , Lampert,	.	.					
<i>Stereoderma murrayi</i> , Bell,	.	.	x				
<i>validum</i> , Bell,	.	.			x		
<i>Thyonidium cebuense</i> , Semper,	.	.			x		
<i>magnum</i> , Ludwig,	.	.			x		
<i>caudatum</i> , Hutton,	.	.			x	x	
<i>japonicum</i> , v. Marenzeller,	.	.			x		
<i>schmeltzii</i> , Ludwig,	.	.			x	x	
<i>molle</i> , Selenka,	.	.			x		
<i>rugosum</i> , n. sp.,	.	.			x		
<i>Orcula limaconotus</i> , Brandt,	.	.		x			
<i>tenera</i> , Ludwig,	.	.			x		x
<i>hypsiptyga</i> , von Marenzeller,	.	.	x			x	
<i>cucumiformis</i> , Semper,	.	.			x		
<i>Phyllophorus perspicillum</i> , Selenka,	.	.				x	
<i>incompertus</i> , n. sp.,	.	.				x	
<i>ehrenbergii</i> , Selenka,	.	.		x			
<i>frauenfeldi</i> , Ludwig,	.	.		x			
<i>proteus</i> , Bell,	.	.		x			
<i>gracilis</i> , Selenka,	.	.		x			
<i>tenuis</i> , Haacke,	.	.	x				
<i>Psolus ornatus</i> , Verrill,	.	.					x
<i>complanatus</i> , Semper,	.	.			x		
<i>boholensis</i> , var. <i>pandanensis</i> , Semper,	.	.			x		
<i>boholensis</i> , Semper,	.	.			x		
<i>fabricii</i> , Düben and Koren,	.	.			x		
<i>cataphractus</i> , Selenka,	.	.				x	
<i>ambulator</i> , Bell,	.	.			x		
<i>appendiculatus</i> , Blainville,	.	.	x				
<i>antarcticus</i> , Philippi,	.	.				x	
<i>squamatus</i> , Düben and Koren,	.	.					x
<i>Labidodemas semperianum</i> , Selenka,	.	.					x