

	PAGE		PAGE
<i>chronhjelmi</i> , n. sp. (Cucumaria), . . .	105	<i>Cuvieria</i> , Péron, Cuvier, . . .	110, 127-131
<i>cigaro</i> , Troschel (Anaperus), . . .	140	<i>cuvieria</i> , Jæger (Cuvieria), . . .	131
<i>cigaro</i> , Selenka (Stolus), . . .	140	<i>cuvieria</i> , Cuvier (Holothuria), . . .	131
<i>cigaro</i> , Selenka (Thyone), . . .	140	<i>cuvierius</i> , Semper (Psolus), . . .	131
<i>cinerascens</i> , Brandt (Stichopus, subg. Gymnochirota), . . .	212	<i>cylindrica</i> , Semper (Cucumaria), . . .	101
<i>cinerascens</i> , Grube (Stichopus), . . .	238	<i>cylindricus</i> , Semper (Colochirus), . . .	122
<i>citrea</i> , Semper (Cucumaria), . . .	104	<i>cylindricus</i> , Haacke (Stichopus), . . .	189
<i>Cladodactyla</i> , Brandt, 100, 101, 102, 103, 104, 110, 113, 125		<i>Cystipus</i> , Haacke, . . .	230
<i>Cladolabes</i> , Brandt, . . .	120, 141, 149	<i>Dactylota</i> , Brandt, . . .	24, 34, 115
<i>clemens</i> , Ludwig (Holothuria), . . .	204	<i>danielsseni</i> , n. sp. (Ankyroderma), . . .	39
<i>cæruleus</i> , Semper (Colochirus), . . .	120	DECACHIROTÆ, Bell (Subfamily), . . .	266
<i>cognita</i> , Lampert (Semperia), . . .	266	<i>decollata</i> , Gray (Holothuria), . . .	116
<i>collaris</i> , Haacke (Holothuria), . . .	227	<i>decorata</i> , von Marenzeller (Holothuria), . . .	218
<i>colloradiatus</i> , Haacke (Colochirus), . . .	124	DENDROCHIROTÆ, Brandt (Family), . . .	57, 99
<i>Colochirus</i> , Troschel, . . .	76, 119	<i>depressa</i> , Ludwig (Holothuria), . . .	213
<i>Colpochirota</i> , Brandt, . . .	202	<i>dicquemarii</i> , Grube (Cladodactyla), . . .	100
<i>coluber</i> , Semper (Holothuria), . . .	232	<i>dicquemarii</i> , Sars (Cucumaria), . . .	100
<i>columnæ</i> , Delle Chiaje (Holothuria), . . .	229	<i>dicquemarii</i> , Cuvier (Holothuria), . . .	106
<i>columnæ</i> , Cuvier (Holothuria), . . .	194	<i>dietrichii</i> , Ludwig (Holothuria), . . .	208
<i>commune</i> , Düben and Koren (Thyonidium), . . .	143	<i>difficilis</i> , Bell (Actinocucumis), . . .	125
<i>communis</i> , Forbes (Cucumaria), . . .	143	<i>difficilis</i> , Semper (Holothuria), . . .	219
<i>complanatus</i> , Semper (Psolus), . . .	127	<i>digitata</i> , Forbes (Chirodota), . . .	29
<i>conchilegum</i> , Pourtalès (Thyonidium), . . .	145	<i>digitata</i> , Lamarck (Fistularia), . . .	29
<i>confundens</i> , n. sp. (Pælopatides), . . .	154	<i>digitata</i> , Montagu (Holothuria), . . .	29
<i>conjungens</i> , Semper (Cucumaria), . . .	112	<i>digitata</i> , Fleming (Mülleria), . . .	29
<i>contorta</i> , Ludwig (Chirodota), . . .	16, 33	<i>digitata</i> , Müller (Synapta), . . .	29
<i>corallifera</i> , Gærtner (Hydra), . . .	106	<i>digitata</i> , Blainville (Thyone), . . .	29
<i>coriacea</i> , Hutton (Caudina), . . .	47, 54	<i>Diploperideris</i> , Brandt, . . .	196
<i>coriacea</i> Hutton (Echinostoma), . . .	54	<i>disciformis</i> , n. sp. (Psolus), . . .	85
<i>coriacea</i> , Hutton (Molpadia), . . .	54	<i>discolor</i> , Eschscholtz (Chirodota), . . .	36
<i>coriacea</i> , Agassiz (Synapta), . . .	34	<i>discolor</i> , n. sp. (Cucumaria), . . .	64
<i>coriaceus</i> , Plancus (Echinus), . . .	101	<i>discrepans</i> , Semper (Holothuria), . . .	226
<i>crocea</i> , Wyville Thomson (Cladodactyla), . . .	110	<i>dissimilis</i> , Leslie and Herdman (Cucumaria), . . .	116
<i>crocea</i> , Selenka (Cucumaria), . . .	110	<i>dissimilis</i> , Fleming (Holothuria), . . .	116
<i>crocea</i> , Lesson (Holothuria), . . .	58, 110	<i>distincta</i> , von Marenzeller (Synapta), . . .	11, 32
<i>crocea</i> , Jæger (Pentacta), . . .	110	<i>dobsoni</i> , Bell (Phyllophorus), . . .	151
<i>crucifera</i> , Semper (Cucumaria), . . .	100	<i>dolabrifera</i> , Stimpson (Synapta), . . .	27
<i>cubana</i> , Ludwig (Holothuria), . . .	237	<i>doliolum</i> , Pallas (Actinia), . . .	125
<i>Cucumaria</i> , Blainville, . . .	57, 99	<i>doliolum</i> , Brandt (Cladodactyla), . . .	125
<i>cucumiformis</i> , Semper (Orcula), . . .	150	<i>doliolum</i> , Grube (Cladodactyla), . . .	101
<i>cucumis</i> , Semper (Colochirus), . . .	82, 123	<i>doliolum</i> , von Marenzeller (Colochirus), . . .	125
<i>cucumis</i> , Sars (Cucumaria), . . .	107	<i>doliolum</i> , Sars (Cucumaria), . . .	101
<i>cucumis</i> , Risso (Holothuria), . . .	107	<i>doliolum</i> , Lamarck (Holothuria), . . .	101, 125
<i>cucumis</i> , Delle Chiaje (Pentacta), . . .	116	<i>doliolum</i> , Jæger (Pentacta), . . .	101, 125
<i>cunninghami</i> , Bell (Thyone), . . .	137	<i>doreyana</i> , Quoy and Gaimard (Fistularia), . . .	23
<i>curiosa</i> , Ludwig (Holothuria), . . .	181, 220	<i>doreyana</i> , Brandt (Synapta), . . .	23
<i>curvata</i> , Lampert (Thyone), . . .	267	<i>drummondii</i> , Forbes (Cucumaria), . . .	143
		<i>drummondii</i> , Thompson (Holothuria), . . .	143
		<i>drummondii</i> , Sars (Thyonidium), . . .	143