

Mitra cineracea, Sowerby, Thes. Conch., vol. iv. pts. 31, 32, p. 32, sp. 438, pl. cclxxiii. (xxii. Gen.)
figs. 494, 495 (*Mitra judæorum*, Dohrn, var.)

Turricula (Costellaria) cineracea, Tryon, Manual, vol. iv. p. 175, pl. lii. fig. 492.

Station 162. April 2, 1874. Lat. 39° 10' 30" S., long. 146° 37' E. Off East Moncœur Island, Bass Strait. 38 fathoms. Sand and shells.

Habitat.—Mindanao, Philippines (Reeve).

18. *Mitra (Pusia) nodulifera*, A. Adams.

Mitra nodulifera, A. Adams, Proc. Zool. Soc. Lond., 1851, p. 140, sp. 48.

„ „ Sowerby, Thes. Conch., vol. iv. pts. 31, 32, p. 32, sp. 436, pl. cclxxv. (xxiv. Gen.)
fig. 530.

„ (*Turricula) nodulifera*, E. A. Smith, Mar. Shells Solomon Islands, Journ. Linn. Soc. Lond.,
vol. xii. p. 550, No. 53.

Turricula (Costellaria) nodulifera, Tryon, Manual, vol. iv. p. 170, pl. l. fig. 442.

July 29, 1874. Levuka, Fiji. 12 fathoms.

Habitat.—?—(Adams); the Solomon Islands and Philippines (E. A. Smith).

19. *Mitra (Callithea) obeliscus*, Reeve.

Mitra obeliscus, Reeve, Proc. Zool. Soc. Lond., 1844, p. 175.

„ „ Reeve, Conch. Icon., vol. ii. pl. xv. fig. 107.

„ „ Sowerby, Thes. Conch., vol. iv. pts. 31, 32, p. 36, sp. 486, pl. cclxx. (ix. Gen.) figs.
126, 127.

„ (*Turricula) obeliscus*, E. A. Smith, Mar. Shells Solomon Islands, Journ. Linn. Soc. Lond.,
vol. xii. p. 549, No. 47.

„ „ „ Brazier, "Chevert" Exped. Shells, Proc. Linn. Soc. New South Wales,
vol. i. p. 214.

Turricula obeliscus, v. Martens, Moll. Mauritius, &c., p. 254.

Mitra (Callithea) obeliscus, E. A. Smith, Andaman Shells, Proc. Zool. Soc. Lond., 1878, p. 815, No. 47.

Turricula (Costularia) obeliscus, Tryon, Manual, vol. iv. p. 179, pl. liii. fig. 535.

July 29, 1874. Levuka, Fiji. 12 fathoms.

Habitat.—Negros, Philippines (Reeve); Torres Strait, 11 to 30 fathoms (Brazier); Solomon and Andaman Islands (E. A. Smith); Mauritius and Indian Ocean (v. Martens).

This species is not unfrequently confounded with *Mitra delicata*, A. Adams, but is markedly more slender, has the pillar lip longer, thinner, and more twisted; the teeth on the outer lip, too, are much deeper within the throat.

20. *Mitra (Callithea) granosa* (Chemnitz).

Chemnitz, Conch. Cab., vol. iv. p. 222, pl. cxlviii. figs. 1373, 1374 (not figs. 1367, 1368).

Voluta granosa, Chemnitz, Conch. Cab., vol. x. pp. 139, 173, pl. cli. figs. 1442, 1443.