

11. *Strombus (Gallinula) variabilis*, Swainson.

- Strombus variabilis*, Swainson, Zool. Illustr., ser. 1, vol. i. pl. x.
 " " Kiener, Iconog., p. 49, sp. 38, pl. xxi. fig. 2.
 " " Sowerby, Genera, vol. ii. pl. ccxxxi. fig. 5.
 " " Sowerby, Thes. Conch., vol. i. pt. 1, p. 27, sp. 9, pl. vi. figs. 9, 13, 14.
 " " Reeve, Conch. Icon., vol. vi. pl. x. fig. 21.
 " " Wood (ed. Hanley), p. 214, suppl. pl. iv. fig. 2.
 " " Küster, Conch. Cab. (ed. Küster), p. 37, sp. 22, pl. iv.a. figs. 1, 2.
Gallinula variabilis, Brazier, "Chevert" Exped. Shells, Proc. Linn. Soc. New South Wales, 1877, vol. i. p. 292, No. 46.

March 7, 1875. Admiralty Islands, north-east of Papua. 16 to 25 fathoms.

Habitat.—North Australia.

12. *Strombus (Gallinula) vittatus*, Linne.

- Strombus vittatus*, Linne, Syst. Nat. (ed. 12), p. 1211, No. 508.
 Martini, Conch. Cab., vol. iii. p. 109, pl. lxxix. figs. 819, 820, 822, 823.
 " " Chemnitz, Conch. Cab., vol. x. p. 208, pl. clv. figs. 1481, 1482.
 " " Dillwyn, Cat., vol. ii. p. 671, sp. 28.
 " " Wood, Ind. Test., p. 124, pl. xxiv. fig. 28 (?).
 " " Lamarck, Anim. s. vert., vol. vii. p. 207, and (ed. Desh.) vol. ix. p. 701, sp. 19.
 " *turritus*, Lamarck, Anim. s. vert., vol. vii. p. 212, and (ed. Desh., see his note) vol. ix. p. 709, sp. 31.
 " *vittatus*, Deshayes, Encycl. method., vers, vol. iii. p. 991, sp. 10, pl. ccccix. fig. 1 (reversed).
 " " Sowerby, Thes. Conch., vol. i. pt. 1, p. 26, sp. 4, pl. vi. figs. 27-31.
 " " Kiener, Iconog., p. 40, sp. 31, pl. xxiii. fig. 1, and (*Strombus turritus*) p. 42, sp. 32, pl. xxiv. fig. 1.
 " " Küster, Conch. Cab. (ed. Küster), p. 44, sp. 29, pl. vii. figs. 5, 6, 8, 9, and (*Strombus turritus*) p. 55, sp. 37, pl. x. figs. 3, 4.
 " " Reeve, Conch. Icon., vol. vi. pl. xvii. fig. 44.
 " " Hanley, Ipsa Linn. Conch., pp. 273, 518.
Gallinula vittata, Brazier, "Chevert" Exped. Shells, Proc. Linn. Soc. New South Wales, 1877, vol. i. p. 293, No. 49.

July 29, 1874. Levuka, Fiji. 12 fathoms.

Station 203. October 31, 1874. Lat. 11° 6' N., long. 123° 9' E. Philippines. 20 fathoms. Mud.

Habitat.—East Indies (Sowerby); Philippines (Reeve and British Museum), also Moluccas (Berne Museum, Collection Shuttleworth); North-east Australia, 15 fathoms (Brazier).

I have marked the reference to Wood (*loc. cit. supra*) with a query. Mr Hanley, in his edition of the Index, accepts Wood's identification of the species, and I daresay he knows best, but I should have thought that Wood's figure characteristically represented *Strombus campbellii*, Gray.