

it is slightly hollowed backward on the edge, while it advances in front to the periphery : its direction is very straight in the line of the spire's slope ; on the base, which is narrow, it is extremely patulous. *Inner lip* : a very thin and undefined glaze expands on the body, which is very faintly convex, and melts without a break into the pillar, which is very slightly thickened, rounded, and twisted, and behind which lies the slightest possible furrow. H. 0·17 in. B. 0·04. Penultimate whorl, height 0·027. Tip of apex, breadth 0·005. Mouth, height 0·058, breadth 0·024.

This species, in the perfect flatness of its whorls, the absence of any impression of the suture, and form of the apex, greatly resembles *Eulima subulata*, Don. ; but it is much more attenuated, the body-whorl in particular is more contracted at the periphery, and the apex is smaller and more perfectly rounded and very slightly bent to one side. In *Eulima bilineata*, Alder, the base of which is more like the Challenger species, the whorls are slightly rounded. In *Eulima philippii*, Wein., var. *gracilis*, there is always some slight irregularity in the spire, and the apex is very much more minute.

5. *Eulima fasciata*, Watson (Pl. XXXV. fig. 2).

*Eulima fasciata*, Watson, Prelim. Report, pt. 17, Journ. Linn. Soc. Lond., vol. xvii. p. 113, sp. 2.

Station 24. March 25, 1873. Lat. 18° 38' 30" N., long. 65° 5' 30" W. Off Culebra Island, West Indies. 390 fathoms. Pteropod ooze.

*Shell*.—Very small, slightly bent, short ; the suture is discernible on the surface and faintly banded above with white, the base is rounded, the mouth pointedly oval, the tip small and rounded. *Sculpture* : none. *Colour* translucent, with an ivory band encircling the body-whorl at the periphery, and lying above the suture on the spire. *Apex* small, perfectly rounded. *Spire* rather attenuated and bent. *Whorls* 9, flat, the last rather long, with a produced rounded base. *Suture* oblique, slightly impressed, especially near the apex. *Mouth* rather large, oval, pointed above. *Outer lip* a little prominently arched ; on its front edge it is rather deeply sinuated above, prominent in the middle, and retreating at the base so as to form a broad, shallow, little canal in front. *Inner lip* : a very narrow rather thin glaze with a defined edge crosses the body, and runs straight down the pillar, which is very narrow and patulous, with a sharp prominent edge and a very minute furrow behind it. H. 0·096 in. B. 0·034. Penultimate whorl, height 0·017. Tip of apex, breadth 0·004. Mouth, height 0·03, breadth 0·017.

Than *Eulima jeffreysiana*, Brus., this is a slightly stamper form ; the whorls are not in the least convex, the apex is blunter and more truncated, the outer lip is much more laterally expanded, and the basal line much more oblique. The form of the mouth and the slightly larger tip distinguish it from the young of *Eulima intermedia*, Cantr.