

22. *Eulima amblic*,<sup>1</sup> Watson (Pl. XXXVII. fig. 4).

*Eulima amblic*, Watson, Prelim. Report, pt. 17, Journ. Linn. Soc. Lond., vol. xvii. p. 127, sp. 20.

Station 145. December 27, 1873. Lat. 46° 43' S., long. 38° 4' 30" E. Half-way between Marion Island and Prince Edward Island. 50 to 150 fathoms. Volcanic sand.

*Shell*.—Small, thin, translucent or transparent, a little cylindrical, slightly bent, with a blunt tip, nearly flat-sided whorls, a very slight linear suture, and an oblong mouth. *Sculpture*: There are some very obscure microscopic lines of growth and spiral scratches. *Colour* glassy, weathering to translucent. *Spire* narrow, slightly unsymmetrical, being a little bent to the right. *Whorls* 6, short, flattened, and scarcely convex on the side; the last is somewhat long with a very blunt round base. *Suture* very slight indeed. *Mouth* oval, pointed above. *Outer lip* curved a little flatly above, but very roundly below; its edge retreats very much above and below, and is prominent at the periphery. *Inner lip* forms a continuous very slightly concave curve across the body and down the short pillar, at the point of which it is slightly angulated at its junction with the base. H. 0·125 in. B. 0·047. Penultimate whorl, height 0·02. Tip of apex, breadth 0·008. Mouth, height 0·04, breadth 0·026.

This species, which is represented by only one specimen and a fragment, is probably full-grown. Compared to *Eulima philippii*, Wein.k, this is a narrower shell, with a much blunter apex, a longer base, a smaller mouth, and an outer lip less laterally prominent.

23. *Eulima hebes*, Watson (Pl. XXXVII. fig. 6).

*Eulima hebes*, Watson, Prelim. Report, pt. 17, Journ. Linn. Soc. Lond., vol. xvii. p. 127, sp. 21.

Station 122. September 10, 1873. Lat. 9° 5' S., long. 34° 50' W. Off Pernambuco. 350 fathoms. Red mud.

*Shell*.—Rather small, broad, blunt, slightly angulated at the periphery, straight, with conical profiles, flat-sided whorls, a slightly impressed suture, and a small angulated mouth. *Sculpture*: There are unequal, close-set, hair-like microscopic lines of growth, and faint broader spirals. *Colour* glossy ivory-white. *Apex* very blunt, rounded, with a slight projection of the extreme tip on one side. *Spire* short, conical, very slightly contracted toward the apex. *Whorls* 6, of regular increase; the last is subangulated at the periphery. *Suture* somewhat oblique, slightly impressed and distinct. *Mouth* somewhat lozenge-shaped, being pointed above and at the end of the pillar, and also at the periphery of the shell and the base of the pillar. *Outer lip* thin, angulated at its edge, retreats above and in front, and is prominent at the periphery. *Inner lip*: a thinnish

<sup>1</sup> ἀμβλίς, blunt.