

12. *Cylichna labiata*, Watson (Pl. L. fig. 4).

*Cylichna labiata*, Watson, Prelim. Report, pt. 19, Journ. Linn. Soc. Lond., vol. xvii. p. 324.

October 6, 1874. Amboina. 15 to 25 fathoms.

*Shell*.—Small, strong, oval, obliquely truncate above, where the apex is impressed, bluntly pointed in front, obscurely angulated above the middle, umbilicate, finely and closely striate, with a thickened lip posteriorly produced. *Sculpture*: Longitudinals—there are fine, straight, hair-like unequal lines of growth. Spirals—there are fine spiral lines, which at the top and bottom of the shell are a little strong and remote, but in all the centre part are very faint and crowded, and above the middle, where a very blunt angulation occurs, they are nearly invisible. *Colour* translucent white, with a tinge of brown, which on the outer lip and base is ochreous. *Mouth* long, curved, narrow, a little enlarged in front. *Outer lip* rises perpendicularly from the outer edge of the impressed apex, is produced posteriorly, where it bends in toward the apex, and is patulous and almost emarginate; its direction and edge line are both somewhat flexuous; in front it is contracted in towards the pillar, forming at the point of the base a narrow, patulous, but not emarginate, gutter; the edge is sharp, but is strengthened a little way within by a small longitudinal rib, which is of a ruddy white colour. *Top* obliquely truncate, small, with a blunt keel round the edge of the minute, shallow, conical depression, which is half choked by the outer lip. *Inner lip*: a very thin glaze extends across the gibbously and convexly curved body; there is a slight angle at the top of the pillar, which is direct, a little oblique, hardly patulous, scarcely twisted, very narrow, most feebly toothed and truncate in front; behind the sharp edge is a small but deep umbilicus. H. 0·16 in. B. 0·09. Mouth, breadth at same place, 0·01.

This is a very peculiar form, extremely like an *Ovula* both in shape and in the thickness of the outer lip, the edge of which, however, is sharp.

13. *Cylichna (Volvula) paupercula*, Watson (Pl. L. fig. 5).

*Cylichna (Volvula) paupercula*, Watson, Prelim. Report, pt. 19, Journ. Linn. Soc. Lond., vol. xvii. p. 325.

Station 24. March 25, 1873. Lat. 18° 38' 30" N., long. 65° 5' 30" W. North of Culebra Island, West Indies. 390 fathoms. Pteropod ooze.

*Shell*.—Small, cylindrically oblong, with a short, blunt, but pointed top, white, faintly spiralled. *Sculpture*: Longitudinals—there are fine close-set lines of growth. Spirals—the whole shell is scored with fine, shallow, remote, scarcely fretted furrows. *Colour* ivory-white. *Mouth* arched, narrow above and throughout the greater part of its length, but widening in front, where the body of the shell contracts on the base; above, it rises