

August 1874. Kandavu, Fiji. Inhabited by a land-crab.

*Habitat*.—From the Natal coast (Krauss) and the Red Sea (Issel), across the whole Indian Ocean to Polynesia and Australia (v. Martens).

Sowerby (Thes. Conch., pt. 40, p. 112, sp. 41, pl. cccclxiv. figs. 26–28, pl. cccclxviii. fig. 118) is not quoted above, as his fig. 27 does not represent this species. The others perhaps do, but it is impossible to be sure.

No. IV. (See p. 132.)

3. *Nerita chamaeleon*, Linne.

*Nerita chamaeleon*, Linne, Syst. Nat. (ed. 12), p. 1255, No. 737.

Chemnitz, Conch. Cab., vol. v. p. 310, pl. cxci. figs. 1988, 1989 (1990, 1991, excl.).

- „ „ Dillwyn, Catalogue, vol. ii. p. 1003, sp. 57.
- „ „ Wood, Ind. Test., p. 181, pl. xxxvi. fig. 60, and (*Nerita stella*) fig. 61.
- „ „ Lamarck, Anim. s. vert., vol. vi. (2) p. 192, sp. 9.
- „ „ Deshayes, Encycl. méthod., vers, vol. iii. p. 618, sp. 4, pl. cccliv. fig. 3 (as *Nerita bizonalis*).
- „ „ Hanley, Ipsa Linn. Conch., pp. 407, 543.
- „ *stella*, Reeve, Conch. Icon., vol. ix. pl. xv. fig. 64.
- „ *chamaeleo*, v. Martens, Moll. Mauritius, &c., p. 291.
- „ *chamaeleon*, Sowerby, Thes. Conch., pt. 40, p. 111, sp. 38, pl. ccclxiv. (ii. Gen.) fig. 22, pl. ccclxvii. (v. Gen.) figs. 94–99.

February 1, 1875. Samboangan, Philippines. 10 fathoms.

*Habitat*.—Madagascar and the Indian Ocean, to the Malay Archipelago (v. Martens).

No. V. (See p. 172.)

3. *Purpura (Thalessa) pica*, de Blainville.

Martini, Conch. Cab., vol. iii. p. 269, pl. c. figs. 956–958.

*Purpura pica*, de Blainville, Pourpre, Nouv. Ann. du Mus., vol. i. p. 213, sp. 27, pl. ix. fig. 9.

- „ *armigera*, var., Quoy and Gaimard, "Astrolabe," Zool., vol. iii. p. 556, pl. xxxvii. figs. 17, 18.
- „ *pica*, Kiener, Iconog., p. 48, sp. 29, pl. xi. fig. 31.
- „ „ Lamarck, Anim. s. vert. (ed. Desh.), vol. x. p. 110, sp. 77.
- „ „ Küster, Conch. Cab. (ed. Küster), p. 190, sp. 98, pl. xxxi. figs. 9, 10.
- „ „ v. Martens, Moll. Mauritius, &c., p. 236.

August 1874. Kandavu, Fiji.

*Habitat*.—Manilla (Küster); from the Malayan Archipelago, across the whole Indian Ocean, to the east coast of Africa (v. Martens).

The solitary specimen of this species was picked up some way inland occupied by a hermit crab (*Cænobia rugosa*, M.-Edw.).